

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## 1. NAME OF THE MEDICINAL PRODUCT

Triumeq 50 mg/600 mg/300 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 50 mg dolutegravir (as sodium), 600 mg of abacavir (as sulfate) and 300 mg of lamivudine.

For the full list of excipients see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet (tablet)

Purple, biconvex, film-coated oval tablets, approximately 22 x 11 mm, debossed with "572 Tr" on one side.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Triumeq is indicated for the treatment of Human Immunodeficiency Virus (HIV) infected adults and adolescents above 12 years of age weighing at least 40 kg (see sections 4.4 and 5.1).

Before initiating treatment with abacavir-containing products, screening for carriage of the HLA-B\*5701 allele should be performed in any HIV-infected patient, irrespective of racial origin (see section 4.4). Abacavir should not be used in patients known to carry the HLA-B\*5701 allele.

### 4.2 Posology and method of administration

Therapy should be prescribed by a physician experienced in the management of HIV infection.

#### Posology

*Adults and adolescents (weighing at least 40kg)*

The recommended dose of Triumeq in adults and adolescents is one tablet once daily.

Triumeq should not be administered to adults or adolescents who weigh less than 40 kg because it is a fixed-dose tablet that cannot be dose reduced.

Triumeq is a fixed-dose tablet and should not be prescribed for patients requiring dose adjustments. Separate preparations of dolutegravir, abacavir or lamivudine are available in cases where discontinuation or dose adjustment of one of the active substances is indicated. In these cases the physician should refer to the individual product information for these medicinal products.

#### *Missed doses*

If the patient misses a dose of Triumeq, the patient should take Triumeq as soon as possible, providing the next dose is not due within 4 hours. If the next dose is due within 4 hours, the patient should not take the missed dose and simply resume the usual dosing schedule.

#### *Elderly*

There are limited data available on the use of dolutegravir, abacavir and lamivudine in patients aged 65 years and over. There is no evidence that elderly patients require a different dose than younger adult patients (see section 5.2). Special care is advised in this age group due to age associated changes such as the decrease in renal function and alteration of haematological parameters.

#### *Renal impairment*

Triumeq is not recommended for use in patients with a creatinine clearance < 50 ml/min (see section 5.2).

#### *Hepatic impairment*

A dose reduction of abacavir may be required for patients with mild hepatic impairment (Child-Pugh grade A). As dose reduction is not possible with Triumeq, the separate preparations of dolutegravir, abacavir or lamivudine should be used when this is judged necessary. Triumeq is not recommended in patients with moderate and severe hepatic impairment (see sections 4.4 and 5.2).

#### *Paediatric population*

The safety and efficacy of Triumeq in children less than 12 years of age has not yet been established. No data are available.

### Method of administration

#### Oral use

Triumeq can be taken with or without food (see section 5.2).

### **4.3 Contraindications**

Hypersensitivity to dolutegravir, abacavir or lamivudine or to any of the excipients listed in section 6.1. See sections 4.4 and 4.8.

Co-administration with dofetilide (see section 4.5).

### **4.4 Special warnings and precautions for use**

#### Transmission of HIV

While effective viral suppression with antiretroviral therapy has been proven to substantially reduce the risk of sexual transmission, a residual risk cannot be excluded. Precautions to prevent transmission should be taken in accordance with national guidelines.

#### Hypersensitivity reactions (see section 4.8)

Both abacavir and dolutegravir are associated with a risk for hypersensitivity reactions (HSR) (see section 4.8), and share some common features such as fever and/or rash with other symptoms indicating multi-organ involvement. Clinically it is not possible to determine whether a HSR with Triumeq would be caused by abacavir or dolutegravir. Hypersensitivity reactions have been observed more commonly with abacavir, some of which have been life-threatening, and in rare cases fatal,

when not managed appropriately. The risk for abacavir HSR to occur is high for patients who test positive for the HLA-B\*5701 allele. However, abacavir HSRs have been reported at a low frequency in patients who do not carry this allele.

Therefore, the following should always be adhered to:

- HLA-B\*5701 status must always be documented prior to initiating therapy.
- Triumeq should never be initiated in patients with a positive HLA-B\*5701 status, nor in patients with a negative HLA-B\*5701 status who had a suspected abacavir HSR on a previous abacavir-containing regimen.
- **Triumeq must be stopped without delay**, even in the absence of the HLA-B\*5701 allele, if an HSR is suspected. Delay in stopping treatment with Triumeq after the onset of hypersensitivity may result in an immediate and life-threatening reaction. Clinical status including liver aminotransferases and bilirubin should be monitored.
- After stopping treatment with Triumeq for reasons of a suspected HSR, **Triumeq or any other medicinal product containing abacavir or dolutegravir must never be re-initiated**.
- Restarting abacavir containing products following a suspected abacavir HSR can result in a prompt return of symptoms within hours. This recurrence is usually more severe than on initial presentation, and may include life-threatening hypotension and death.
- In order to avoid restarting abacavir and dolutegravir, patients who have experienced a suspected HSR should be instructed to dispose of their remaining Triumeq tablets.

#### Clinical description of HSRs

Hypersensitivity reactions have been reported in <1% of patients treated with dolutegravir in clinical studies, and were characterized by rash, constitutional findings, and sometimes, organ dysfunction, including severe liver reactions.

Abacavir HSR has been well characterised through clinical studies and during post marketing follow-up. Symptoms usually appeared within the first six weeks (median time to onset 11 days) of initiation of treatment with abacavir, **although these reactions may occur at any time during therapy**.

Almost all HSR to abacavir will include fever and/or rash. Other signs and symptoms that have been observed as part of abacavir HSR are described in detail in section 4.8 (Description of selected adverse reactions), including respiratory and gastrointestinal symptoms. Importantly, such symptoms **may lead to misdiagnosis of HSR as respiratory disease (pneumonia, bronchitis, pharyngitis), or gastroenteritis**. The symptoms related to this HSR worsen with continued therapy and **can be life-threatening**. These symptoms usually resolve upon discontinuation of abacavir.

Rarely, patients who have stopped abacavir for reasons other than symptoms of HSR have also experienced life-threatening reactions within hours of re- initiating abacavir therapy (see Section 4.8 Description of selected adverse reactions). Restarting abacavir in such patients must be done in a setting where medical assistance is readily available.

### Lactic acidosis

Lactic acidosis, usually associated with hepatomegaly and hepatic steatosis, has been reported with the use of nucleoside analogues. Early symptoms (symptomatic hyperlactataemia) include benign digestive symptoms (nausea, vomiting and abdominal pain), non-specific malaise, loss of appetite, weight loss, respiratory symptoms (rapid and/or deep breathing) or neurological symptoms (including motor weakness).

Lactic acidosis has a high mortality and may be associated with pancreatitis, liver failure, or renal failure.

Lactic acidosis generally occurred after a few or several months of treatment.

Treatment with nucleoside analogues should be discontinued in the setting of symptomatic hyperlactataemia and metabolic/lactic acidosis, progressive hepatomegaly, or rapidly elevating aminotransferase levels.

Caution should be exercised when administering nucleoside analogues to any patient (particularly obese women) with hepatomegaly, hepatitis or other known risk factors for liver disease and hepatic steatosis (including certain medicinal products and alcohol). Patients co-infected with hepatitis C and treated with alpha interferon and ribavirin may constitute a special risk.

Patients at increased risk should be followed closely.

### Lipodystrophy

Combination antiretroviral therapy has been associated with the redistribution of body fat (lipodystrophy) in HIV patients. The long-term consequences of these events are currently unknown. Knowledge about the mechanism is incomplete. A connection between visceral lipomatosis and protease inhibitors (PIs) and lipoatrophy and nucleoside reverse transcriptase inhibitors (NRTIs) has been hypothesised. A higher risk of lipodystrophy has been associated with individual factors such as older age, and with drug related factors such as longer duration of antiretroviral treatment and associated metabolic disturbances. Clinical examination should include evaluation for physical signs of fat redistribution. Consideration should be given to the measurement of fasting serum lipids and blood glucose. Lipid disorders should be managed as clinically appropriate (see section 4.8).

### Liver disease

The safety and efficacy of Triumeq has not been established in patients with significant underlying liver disorders. Triumeq is not recommended in patients with moderate to severe hepatic impairment (see section 4.2).

Patients with pre-existing liver dysfunction, including chronic active hepatitis have an increased frequency of liver function abnormalities during combination antiretroviral therapy, and should be monitored according to standard practice. If there is evidence of worsening liver disease in such patients, interruption or discontinuation of treatment must be considered.

### Patients with chronic hepatitis B or C

Patients with chronic hepatitis B or C and treated with combination antiretroviral therapy are at an increased risk of severe and potentially fatal hepatic adverse reactions. In case of concomitant

antiviral therapy for hepatitis B or C, please refer also to the relevant product information for these medicinal products.

Triumeq includes lamivudine, which is active against hepatitis B. Abacavir and dolutegravir lack such activity. Lamivudine monotherapy is generally not considered an adequate treatment for hepatitis B, since the risk for hepatitis B resistance development is high. If Triumeq is used in patients co-infected with hepatitis B an additional antiviral is therefore generally needed. Reference should be made to treatment guidelines.

If Triumeq is discontinued in patients co-infected with hepatitis B virus, periodic monitoring of both liver function tests and markers of HBV replication is recommended, as withdrawal of lamivudine may result in an acute exacerbation of hepatitis.

As abacavir and ribavirin share the same phosphorylation pathways, a possible intracellular interaction between these medicinal products has been postulated, which could lead to a reduction in intracellular phosphorylated metabolites of ribavirin and, as a possible consequence, a reduced chance of sustained virological response (SVR) for hepatitis C (HCV) in HCV co-infected patients treated with pegylated interferon plus ribavirin. Conflicting clinical findings are reported in literature on co-administration between abacavir and ribavirin. Some data suggest that HIV/HCV co-infected patients receiving abacavir-containing ART may be at risk of a lower response rate to pegylated interferon/ribavirin therapy. Caution should be exercised when medicinal products containing abacavir and ribavirin are co-administered (see section 4.5).

#### Immune Reactivation Syndrome

In HIV-infected patients with severe immune deficiency at the time of institution of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic pathogens may arise and cause serious clinical conditions, or aggravation of symptoms. Typically, such reactions have been observed within the first few weeks or months of initiation of CART. Relevant examples are cytomegalovirus retinitis, generalised and/or focal mycobacterial infections, and *Pneumocystis carinii* pneumonia. Any inflammatory symptoms should be evaluated and treatment instituted when necessary. Autoimmune disorders (such as Graves' disease) have also been reported to occur in the setting of immune reactivation; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment.

Liver chemistry elevations consistent with immune reconstitution syndrome were observed in some hepatitis B and/or C co-infected patients at the start of dolutegravir therapy. Monitoring of liver chemistries is recommended in patients with hepatitis B and/or C co-infection. (See 'Patients with chronic hepatitis B or C' earlier in this section and also see section 4.8).

#### Mitochondrial dysfunction

Nucleoside and nucleotide analogues have been demonstrated *in vitro* and *in vivo* to cause a variable degree of mitochondrial damage. There have been reports of mitochondrial dysfunction in HIV-negative infants exposed *in utero* and/or post-natally to nucleoside analogues. The main adverse reactions reported are haematological disorders (anaemia, neutropenia), metabolic disorders (hyperlactataemia, hyperlipasemia). These reactions are often transitory. Some late-onset neurological disorders have been reported (hypertonia, convulsion, abnormal behaviour). Whether the neurological disorders are transient or permanent is currently unknown. Any child exposed *in utero* to nucleoside and nucleotide analogues, even HIV-negative children, should have clinical and laboratory follow-up and should be fully investigated for possible mitochondrial dysfunction in case of relevant signs or symptoms. These findings do not affect current national recommendations to use antiretroviral therapy in pregnant women to prevent vertical transmission of HIV.

### Myocardial infarction

Observational studies have shown an association between myocardial infarction and the use of abacavir. Those studied were mainly antiretroviral experienced patients. Data from clinical trials showed limited numbers of myocardial infarction and could not exclude a small increase in risk. Overall the available data from observational cohorts and from randomised trials show some inconsistency so can neither confirm nor refute a causal relationship between abacavir treatment and the risk of myocardial infarction. To date, there is no established biological mechanism to explain a potential increase in risk. When prescribing Triumeq, action should be taken to try to minimize all modifiable risk factors (e.g. smoking, hypertension, and hyperlipidaemia).

### Osteonecrosis

Although the aetiology is considered to be multifactorial (including corticosteroid use, bisphosphonates, alcohol consumption, severe immunosuppression, higher body mass index), cases of osteonecrosis have been reported particularly in patients with advanced HIV-disease and/or long-term exposure to CART. Patients should be advised to seek medical advice if they experience joint aches and pain, joint stiffness or difficulty in movement.

### Opportunistic infections

Patients should be advised that Triumeq or any other antiretroviral therapy does not cure HIV infection and that they may still develop opportunistic infections and other complications of HIV infection. Therefore, patients should remain under close clinical observation by physicians experienced in the treatment of these associated HIV diseases.

### Drug resistance

Since the recommended dose of dolutegravir is 50 mg twice daily for patients with resistance to integrase inhibitors, the use of Triumeq is not recommended for patients with integrase inhibitor resistance.

### Drug interactions

Since the recommended dose of dolutegravir is 50 mg twice daily when co-administered with etravirine (without boosted protease inhibitors), efavirenz, nevirapine, rifampicin, tipranavir/ritonavir, carbamazepine, phenytoin, phenobarbital and St. John's wort, the use of Triumeq is not recommended for patients taking these medicines (see section 4.5).

Triumeq should not be co-administered with polyvalent cation-containing antacids. Triumeq is recommended to be administered 2 hours before or 6 hours after these agents (see section 4.5).

Triumeq is recommended to be administered 2 hours before or 6 hours after taking calcium or iron supplements (see section 4.5).

Dolutegravir increased metformin concentrations. A dose adjustment of metformin should be considered when starting and stopping coadministration of dolutegravir with metformin, to maintain glycaemic control (see section 4.5). Metformin is eliminated renally and therefore it is of importance to monitor renal function when co-treated with dolutegravir. This combination may increase the risk for lactic acidosis in patients with moderate renal impairment (stage 3a creatinine clearance [CrCl] 45– 59 mL/min) and a cautious approach is recommended. Reduction of the metformin dose should be highly considered.

The combination of lamivudine with cladribine is not recommended (see section 4.5).

Triumeq should not be taken with any other medicinal products containing dolutegravir, abacavir, lamivudine or emtricitabine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Triumeq contains dolutegravir, abacavir and lamivudine, therefore any interactions identified for these individually are relevant to Triumeq. No clinically significant drug interactions are expected between dolutegravir, abacavir and lamivudine.

##### Effect of other agents on the pharmacokinetics of dolutegravir, abacavir and lamivudine

Dolutegravir is eliminated mainly through metabolism by UGT1A1. Dolutegravir is also a substrate of UGT1A3, UGT1A9, CYP3A4, P-gp, and BCRP. Co-administration of Triumeq and other drugs that inhibit UGT1A1, UGT1A3, UGT1A9, CYP3A4, and/or P-gp may therefore increase dolutegravir plasma concentration. Drugs that induce those enzymes or transporters may decrease dolutegravir plasma concentration and reduce the therapeutic effect of dolutegravir (see Table 1).

The absorption of dolutegravir is reduced by certain anti-acid agents (see Table 1).

Abacavir is metabolised by UDP-glucuronyltransferase (UGT) enzymes and alcohol dehydrogenase; co-administration of inducers or inhibitors of UGT enzymes or with compounds eliminated through alcohol dehydrogenase could alter abacavir exposure.

Lamivudine is cleared renally. Active renal secretion of lamivudine in the urine is mediated through the organic cation transporter (OCT2) and multidrug and toxin extrusion transporters (MATE1 and MATE2-K). Co-administration of lamivudine with OCT and MATE inhibitors could increase lamivudine exposure. Dolutegravir is an OCT2 and MATE1 inhibitor; however, lamivudine concentrations were similar with or without co-administration of dolutegravir based on a cross study analysis, indicating that dolutegravir has no effect on lamivudine exposure *in vivo*.

Abacavir and lamivudine are not significantly metabolised by CYP enzymes.

##### Effect of dolutegravir, abacavir and lamivudine on the pharmacokinetics of other agents

*In vivo*, dolutegravir did not have an effect on midazolam, a CYP3A4 probe. Based on *in vivo* and/or *in vitro* data, dolutegravir is not expected to affect the pharmacokinetics of medicinal products that are substrates of any major enzyme or transporter such as CYP3A4, CYP2C9 and P-gp (for more information see section 5.2).

*In vitro*, dolutegravir inhibited the renal transporters OCT2 and MATE1. *In vivo*, a 10-14% decrease of creatinine clearance (secretory fraction is dependent on OCT2 and MATE-1 transport) was observed in patients. *In vivo*, dolutegravir may increase plasma concentrations of medicinal products in which excretion is dependent upon OCT2 or MATE-1 (e.g. dofetilide, metformin) (see Table 1 and section 4.3).

*In vitro*, dolutegravir inhibited the renal uptake organic anion transporters (OAT)1 and OAT3. Based on the lack of effect on the *in vivo* pharmacokinetics of the OAT substrate tenofovir, *in vivo* inhibition of OAT1 is unlikely. Inhibition of OAT3 has not been studied *in vivo*. Dolutegravir may increase plasma concentrations of medical products in which excretion is dependent upon OAT3.

Abacavir and lamivudine do not inhibit or induce CYP enzymes (such as CYP 3A4, CYP 2C9 or CYP 2D6). *In vitro* data indicate that inhibition of P-pg and BCRP by abacavir cannot be excluded at intestinal level. *In vitro*, lamivudine inhibited OCT1 and OCT2.

Established and theoretical interactions with selected antiretrovirals and non-antiretroviral medicinal products are listed in Table 1.

#### Interaction table

Interactions between dolutegravir, abacavir, lamivudine and co-administered medical products are listed in Table 1 (increase is indicated as “↑”, decrease as “↓”, no change as “↔”, area under the concentration versus time curve as “AUC”, maximum observed concentration as “ $C_{max}$ ”). The table should not be considered exhaustive but is representative of the classes studied.

Table 1: Drug Interactions

Medicinal products by therapeutic areas	Interaction geometric mean change (%)	Recommendations concerning co-administration
<b>Antiretroviral medicinal products</b>		
<i>Non-nucleoside reverse transcriptase inhibitors</i>		
Etravirine without boosted protease inhibitors / Dolutegravir	Dolutegravir ↓ AUC ↓ 71% $C_{max}$ ↓ 52% $C\tau$ ↓ 88%  Etravirine ↔ (induction of UGT1A1 and CYP3A enzymes)	Etravirine without boosted protease inhibitors decreased plasma dolutegravir concentration. Since the recommended dose of dolutegravir is 50 mg twice daily for patients taking etravirine without boosted protease inhibitors, Triumeq is not recommended for patients taking etravirine without co-administration of atazanavir/ritonavir, darunavir/ritonavir or lopinavir/ritonavir (see further below in table).
Lopinavir+ritonavir+etravirine/ Dolutegravir	Dolutegravir ↔ AUC ↑ 11% $C_{max}$ ↑ 7% $C\tau$ ↑ 28%  Lopinavir ↔ Ritonavir ↔ Etravirine ↔	No dose adjustment is necessary.
Darunavir+ritonavir+etravirine/ Dolutegravir	Dolutegravir ↓ AUC ↓ 25% $C_{max}$ ↓ 12% $C\tau$ ↓ 36%  Darunavir ↔ Ritonavir ↔ Etravirine ↔	No dose adjustment is necessary.
Efavirenz/Dolutegravir	Dolutegravir ↓ AUC ↓ 57% $C_{max}$ ↓ 39% $C\tau$ ↓ 75%  Efavirenz ↔ (historical	Since the dose of dolutegravir is 50 mg twice daily when co-administered with efavirenz, the co-administration of efavirenz with Triumeq is not recommended (see section 4.4).

	controls) (induction of UGT1A1 and CYP3A enzymes)	
Nevirapine/Dolutegravir	Dolutegravir ↓ (Not studied, a similar reduction in exposure as observed with efavirenz is expected, due to induction)	Co-administration with nevirapine may decrease dolutegravir plasma concentration due to enzyme induction and has not been studied. Effect of nevirapine on dolutegravir exposure is likely similar to or less than that of efavirenz. Since the dose of dolutegravir is 50 mg twice daily when co-administered with nevirapine, the co-administration of nevirapine with Triumeq is not recommended.
Rilpivirine	Dolutegravir ↔ AUC ↑ 12% $C_{max}$ ↑ 13% $C\tau$ ↑ 22% Rilpivirine ↔	No dose adjustment is necessary.
<i>Nucleoside reverse transcriptase inhibitors (NRTIs)</i>		
Tenofovir  Emtricitabine, didanosine, stavudine, zidovudine.	Dolutegravir ↔ AUC ↑ 1% $C_{max}$ ↓ 3% $C\tau$ ↓ 8% Tenofovir ↔  Interaction not studied	No dose adjustment is necessary when Triumeq is combined with nucleoside reverse transcript inhibitors.  Triumeq is not recommended for use in combination with emtricitabine containing products, since both lamivudine (in Triumeq) and emtricitabine are cytidine analogues (i.e. risk for intracellular interactions, (see section 4.4))
<i>Protease inhibitors</i>		
Atazanavir/Dolutegravir	Dolutegravir ↑ AUC ↑ 91% $C_{max}$ ↑ 50% $C\tau$ ↑ 180%  Atazanavir ↔ (historical controls) (inhibition of UGT1A1 and CYP3A enzymes)	No dose adjustment is necessary.
Atazanavir+ ritonavir/ Dolutegravir	Dolutegravir ↑ AUC ↑ 62% $C_{max}$ ↑ 34% $C\tau$ ↑ 121%  Atazanavir ↔ Ritonavir ↔	No dose adjustment is necessary.
Tipranavir+ritonavir/ Dolutegravir	Dolutegravir ↓ AUC ↓ 59% $C_{max}$ ↓ 47%	Since the recommended dose of dolutegravir is 50 mg twice daily when co-administered with

	$C\tau \downarrow 76\%$ Tipranavir ↔ Ritonavir ↔ (induction of UGT1A1 and CYP3A enzymes)	tipranavir/ritonavir, the co-administration of tipranavir/ritonavir with Triumeq is not recommended.
Fosamprenavir+ritonavir/ Dolutegravir	Dolutegravir↓ $AUC \downarrow 35\%$ $C_{max} \downarrow 24\%$ $C\tau \downarrow 49\%$  Fosamprenavir↔ Ritonavir ↔ (induction of UGT1A1 and CYP3A enzymes)	Fosamprenavir/ritonavir decreases dolutegravir concentrations, but based on limited data, did not result in decreased efficacy in Phase III studies. No dose adjustment is necessary.
Nelfinavir/Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is necessary.
Lopinavir+ritonavir/ Dolutegravir	Dolutegravir ↔ $AUC \downarrow 4\%$ $C_{max} \leftrightarrow 0\%$ $C_{24} \downarrow 6\%$  Lopinavir ↔ Ritonavir ↔	No dose adjustment is necessary.
Darunavir+ritonavir/ Dolutegravir	Dolutegravir ↓ $AUC \downarrow 22\%$ $C_{max} \downarrow 11\%$ $C\tau \downarrow 38\%$  Darunavir ↔ Ritonavir ↔ (induction of UGT1A1 and CYP3A enzymes)	No dose adjustment is necessary.
<b>Other antiviral agents</b>		
Telaprevir	Dolutegravir ↑ $AUC \uparrow 25\%$ $C_{max} \uparrow 19\%$ $C\tau \uparrow 37\%$  Telaprevir ↔ (historical controls) (inhibition of CYP3A enzyme)	No dose adjustment is necessary.
Boceprevir	Dolutegravir ↔ $AUC \uparrow 7\%$ $C_{max} \uparrow 5\%$ $C\tau \uparrow 8\%$  Boceprevir ↔ (historical controls)	No dose adjustment is necessary.
Ribavirin/Abacavir	Interaction not studied. Both drugs are guanosine analogues, and there is a potential to reduce intracellular	Caution should be exercised when both drugs are co-administered (see section 4.4).

	phosphorylated metabolites.	
Daclatasvir/Dolutegravir	Dolutegravir ↔ AUC ↑ 33% $C_{max}$ ↑ 29% $C\tau$ ↑ 45% Daclatasvir ↔	Daclatasvir did not change dolutegravir plasma concentration to a clinically relevant extent. Dolutegravir did not change daclatasvir plasma concentration. No dose adjustment is necessary.
<b>Anti-infective products</b>		
Trimethoprim/sulfamethoxazole (Co-trimoxazole)/Abacavir	Interaction not studied	No Triumeq dosage adjustment necessary, unless patient has renal impairment (See Section 4.2).
Trimethoprim/sulfamethoxazole (Co-trimoxazole)/Lamivudine (160mg/800mg once daily for 5 days/300mg single dose)	Lamivudine: AUC ↑ 43% $C_{max}$ ↑ 7%  Trimethoprim: AUC ↔  Sulfamethoxazole: AUC ↔  (organic cation transporter inhibition)	
<b>Antimycobacterials</b>		
Rifampicin/Dolutegravir	Dolutegravir ↓ AUC ↓ 54% $C_{max}$ ↓ 43% $C\tau$ ↓ 72% (induction of UGT1A1 and CYP3A enzymes)	Since the dose of dolutegravir is 50 mg twice daily when co-administered with rifampicin, the co-administration of rifampicin with Triumeq is not recommended.
Rifabutin	Dolutegravir ↔ AUC ↓ 5% $C_{max}$ ↑ 16% $C\tau$ ↓ 30% (induction of UGT1A1 and CYP3A enzymes)	No dose adjustment is necessary.
<b>Anticonvulsants</b>		
Carbamazepine/Dolutegravir	Dolutegravir ↓ AUC ↓ 49% $C_{max}$ ↓ 33% $C\tau$ ↓ 73%	Since the recommended dose of dolutegravir is 50 mg twice daily when co-administered with carbamazepine, DTG/ABC/3TC FDC is not recommended for patients taking carbamazepine.
Phenobarbital/Dolutegravir Phenytoin/Dolutegravir Oxcarbazepine/Dolutegravir	Dolutegravir ↓ (Not studied, decrease expected due to induction of UGT1A1 and CYP3A enzymes, a similar reduction in exposure as observed with carbamazepine is expected)	Since the recommended dose of dolutegravir is 50 mg twice daily when co-administered with these metabolic inducers, DTG/ABC/3TC FDC is not recommended for patients taking these metabolic inducers.
<b>Antihistamines (histamine H<sub>2</sub> receptor antagonists)</b>		

Ranitidine	Interaction not studied.  Clinically significant interaction unlikely.	No dosage adjustment necessary.
Cimetidine	Interaction not studied.  Clinically significant interaction unlikely.	No dosage adjustment necessary.
<b>Cytotoxics</b>		
Cladribine/Lamivudine	Interaction not studied.  <i>In vitro</i> lamivudine inhibits the intracellular phosphorylation of cladribine leading to a potential risk of cladribine loss of efficacy in case of combination in the clinical setting. Some clinical findings also support a possible interaction between lamivudine and cladribine	Concomitant use of Triumeq with cladribine is not recommended (see section 4.4).
<b>Opioids</b>		
Methadone/Abacavir (40 to 90mg once daily for 14 days/600mg single dose, then 600mg twice daily for 14 days)	Abacavir: AUC ↔ Cmax ↓35%  Methadone: CL/F ↑22%	Methadone dosage adjustment likely not needed in majority of patients; occasionally methadone re-titration may be required.
<b>Retinoids</b>		
Retinoid compounds (e.g. Isotretinoin)	Interaction not studied  Possible interaction given common pathway of elimination via alcohol dehydrogenase (abacavir-component).	Insufficient data to recommend dosage adjustment.
<b>Miscellaneous</b>		
<i>Alcohol</i>		
Ethanol/Dolutegravir Ethanol/Lamivudine	Interaction not studied (Inhibition of alcohol dehydrogenase)	No dosage adjustment necessary.
Ethanol/Abacavir (0.7 g/kg single dose/600mg single dose)	Abacavir: AUC ↑ 41% Ethanol: AUC ↔	
<i>Antiarrhythmics</i>		
Dofetilide/Dolutegravir	Dofetilide ↑ (Not studied, potential increase via inhibition of	Triumeq and dofetilide co-administration is contraindicated due to potential life-threatening toxicity caused

	OCT2 transporter)	by high dofetilide concentration (see section 4.3).
<i>Antacids and supplements</i>		
Magnesium/aluminium-containing antacids/Dolutegravir	Dolutegravir ↓ AUC ↓ 74% $C_{max}$ ↓ 72%  (Complex binding to polyvalent ions)	Magnesium/ aluminium-containing antacids should be taken well separated in time from the administration of Triumeq (minimum 2 hours after or 6 hours before).
Calcium supplements/Dolutegravir	Dolutegravir ↓ AUC ↓ 39% $C_{max}$ ↓ 37% $C_{24}$ ↓ 39%  (Complex binding to polyvalent ions)	Calcium supplements, iron supplements or multivitamins should be taken well separated in time from the administration of Triumeq (minimum 2 hours after or 6 hours before).
Iron supplements/Dolutegravir	Dolutegravir ↓ AUC ↓ 54% $C_{max}$ ↓ 57% $C_{24}$ ↓ 56%  (Complex binding to polyvalent ions)	
Multivitamins/Dolutegravir	Dolutegravir ↓ AUC ↓ 33% $C_{max}$ ↓ 35% $C_{24}$ ↓ 32%	
<i>Corticosteroids</i>		
Prednisone	Dolutegravir ↔ AUC ↑ 11% $C_{max}$ ↑ 6% $C\tau$ ↑ 17%	No dose adjustment is necessary.
<i>Antidiabetics</i>		
Metformin/Dolutegravir	Metformin ↑ Dolutegravir ↔ When co-administered with dolutegravir 50mg QD: Metformin AUC ↑ 79% $C_{max}$ ↑ 66% When co-administered with dolutegravir 50mg BID: Metformin AUC ↑ 145 % $C_{max}$ ↑ 111%	A dose adjustment of metformin should be considered when starting and stopping coadministration of dolutegravir with metformin, to maintain glycaemic control. In patients with moderate renal impairment a dose adjustment of metformin should be considered when coadministered with dolutegravir, because of the increased risk for lactic acidosis in patients with moderate renal impairment due to increased metformin concentration (section 4.4).
<i>Herbal products</i>		
St. John's wort/Dolutegravir	Dolutegravir ↓ (Not studied, decrease expected due to induction of UGT1A1 and CYP3A enzymes, a similar reduction in exposure as	Since the recommended dose of dolutegravir is 50 mg twice daily when co-administered with St. John's wort, DTG/ABC/3TC FDC is not recommended.

	observed with carbamazepine is expected)	
<b><i>Oral contraceptives</i></b>		
Ethinyl estradiol (EE) and Norgestromin (NGMN)/Dolutegravir	Effect of dolutegravir: EE ↔ AUC ↑ 3% $C_{max}$ ↓ 1%  Effect of dolutegravir: NGMN ↔ AUC ↓ 2% $C_{max}$ ↓ 11%	Dolutegravir had no Pharmacodynamic effect on Luteinizing Hormone (LH), Follicle Stimulating Hormone (FSH) and progesterone. No dose adjustment of oral contraceptives is necessary when co-administered with Triumeq.

### Paediatric population

Interaction studies have only been performed in adults.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

As a general rule, when deciding to use antiretroviral agents for the treatment of HIV infection in pregnant women and consequently for reducing the risk of HIV vertical transmission to the newborn, the animal data as well as the clinical experience in pregnant women should be taken into account.

There are no data on the use of Triumeq in pregnancy.

There are no or limited amount of data from the use of dolutegravir in pregnant women. The effect of dolutegravir on human pregnancy is unknown. A moderate amount of data on pregnant women taking the individual actives abacavir and lamivudine in combination indicates no malformative toxicity (more than 400 outcomes from first trimester exposures). Concerning lamivudine, a large amount of data (more than 3000 outcomes from first trimester) indicates no malformative toxicity. Moderate amount of data (more than 600 outcomes from first trimester) indicates no malformative toxicity for abacavir.

In reproductive toxicity studies in animals, dolutegravir was shown to cross the placenta. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). Abacavir and lamivudine may inhibit cellular DNA replication and abacavir has been shown to be carcinogenic in animal models (see section 5.3). The clinical relevance of these findings is unknown.

Triumeq should be used during pregnancy only if the expected benefit justifies the potential risk to the foetus.

For patients co-infected with hepatitis B who are being treated with a lamivudine containing medicinal product such as Triumeq and subsequently become pregnant, consideration should be given to the possibility of a recurrence of hepatitis on discontinuation of lamivudine.

### *Mitochondrial dysfunction*

Nucleoside and nucleotide analogues have been demonstrated *in vitro* and *in vivo* to cause a variable degree of mitochondrial damage. There have been reports of mitochondrial dysfunction in HIV-negative infants exposed *in utero* and/or post-natally to nucleoside analogues (see section 4.4).

### Breast-feeding

It is unknown whether dolutegravir is excreted in human milk. Available toxicological data in animals has shown excretion of dolutegravir in milk. In lactating rats that received a single oral dose of 50 mg/kg at 10 days postpartum, dolutegravir was detected in milk at concentrations typically higher than blood.

Abacavir and its metabolites are excreted into the milk of lactating rats. Abacavir is also excreted into human milk.

Based on more than 200 mother/child pairs treated for HIV, serum concentrations of lamivudine in breastfed infants of mothers treated for HIV are very low (< 4% of maternal serum concentrations) and progressively decrease to undetectable levels when breastfed infants reach 24 weeks of age. There are no data available on the safety of abacavir and lamivudine when administered to babies less than three months old.

It is recommended that HIV infected women do not breast-feed their infants under any circumstances in order to avoid transmission of HIV.

### Fertility

There are no data on the effects of dolutegravir, abacavir or lamivudine on human male or female fertility. Animal studies indicate no effects of dolutegravir, abacavir or lamivudine on male or female fertility (see section 5.3).

## **4.7 Effects on ability to drive and use machines**

Patients should be informed that dizziness has been reported during treatment with dolutegravir. The clinical status of the patient and the adverse reaction profile of Triumeq should be borne in mind when considering the patient's ability to drive or operate machinery.

## **4.8 Undesirable effects**

### Summary of the safety profile

Clinical safety data with Triumeq are limited. The most frequently reported adverse reactions considered possibly or probably related to dolutegravir and abacavir/lamivudine [pooled data from 679 anti retroviral naïve subjects receiving this combination in the Phase IIb to IIIb clinical trials, see section 5.1], were nausea (12%), insomnia (7%), dizziness (6%) and headache (6%).

Many of the adverse reactions listed in the table below occur commonly (nausea, vomiting, diarrhoea, fever, lethargy, rash) in patients with abacavir hypersensitivity. Therefore, patients with any of these symptoms should be carefully evaluated for the presence of this hypersensitivity (see section 4.4). Very rarely cases of erythema multiforme, Stevens-Johnson syndrome or toxic epidermal necrolysis have been reported where abacavir hypersensitivity could not be ruled out. In such cases medicinal products containing abacavir should be permanently discontinued.

The most severe adverse event possibly related to the treatment with dolutegravir and abacavir/lamivudine, seen in individual patients, was a hypersensitivity reaction that included rash and severe liver effects (see section 4.4 and Description of selected adverse reactions in this section).

### Tabulated list of adverse reactions

The adverse reactions considered at least possibly related to treatment with the components of Triumeq from clinical study and post-marketing experience are listed in Table 2 by body system, organ class and absolute frequency. Frequencies are defined as very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1000$ ), very rare ( $< 1/10,000$ ).

The adverse reactions observed for the combination of dolutegravir + abacavir/lamivudine in an analysis of pooled data from Phase IIb to Phase IIIb clinical trials were generally consistent with the adverse reaction profiles for the individual components dolutegravir, abacavir and lamivudine.

There was no difference between the combination and the individual components in severity for any observed adverse reactions.

Table 2: Tabulated summary of adverse reactions associated with the combination of dolutegravir + abacavir/lamivudine in an analysis of pooled data from Phase IIb to Phase IIIb clinical trials and adverse reactions to treatment with abacavir and lamivudine from clinical studies and post-marketing experience, when used with other antiretrovirals

<b>Frequency</b>	<b>Adverse reaction</b>
<i>Blood and lymphatic systems disorders:</i>	
Uncommon:	Neutropenia <sup>2</sup> , anaemia <sup>2</sup> , thrombocytopenia <sup>1</sup>
Very rare:	pure red cell aplasia <sup>1</sup>
<i>Immune system disorders:</i>	
Common	hypersensitivity (see section 4.4) <sup>2</sup>
Uncommon:	immune reconstitution syndrome (see section 4.4) <sup>2</sup>
<i>Metabolism and nutrition disorders:</i>	
Common:	anorexia <sup>1</sup>
Uncommon:	hypertriglyceridaemia, hyperglycaemia
<i>Psychiatric disorders:</i>	
Very common:	insomnia
Common:	abnormal dreams, depression, nightmare, sleep disorder
Uncommon	suicidal ideation or suicide attempt (particularly in patients with a pre-existing history of depression or psychiatric illness)
<i>Nervous system disorders:</i>	
Very common:	headache
Common:	dizziness, somnolence, lethargy <sup>2</sup>
Very rare:	peripheral neuropathy <sup>2</sup> , paraesthesia <sup>2</sup>
<i>Respiratory, thoracic and mediastinal disorders:</i>	
Common:	cough <sup>2</sup> , nasal symptoms <sup>1</sup>
<i>Gastrointestinal disorders:</i>	

Very common:	nausea, diarrhoea
Common:	vomiting, flatulence, abdominal pain <sup>2</sup> , abdominal pain upper <sup>2</sup> , abdominal distension, abdominal discomfort, gastro-oesophageal reflux disease, dyspepsia
Rare:	pancreatitis <sup>2</sup>
<i>Hepatobiliary disorders:</i>	
Uncommon:	hepatitis <sup>2</sup>
<i>Skin and subcutaneous tissue disorders:</i>	
Common:	rash, pruritus, alopecia
Very rare:	erythema multiform <sup>1</sup> , Stevens-Johnson syndrome <sup>1</sup> , toxic epidermal necrolysis <sup>1</sup>
<i>Musculoskeletal and connective tissue disorders:</i>	
Common:	arthralgia <sup>2</sup> , muscle disorders <sup>1</sup>
Rare:	rhabdomyolysis <sup>2</sup>
<i>General disorders and administration site conditions:</i>	
Very common:	fatigue
Common:	asthenia, fever <sup>2</sup> , malaise <sup>2</sup>
<i>Investigations:</i>	
Common:	CPK elevations <sup>2</sup> , ALT/AST elevations <sup>2</sup>
Rare:	amylase elevations <sup>1</sup>
<p><sup>1</sup>This adverse reaction was not identified from the Phase 3 clinical studies for either Triumeq (dolutegravir + abacavir/lamivudine) or dolutegravir, but from clinical studies or post-marketing experience for abacavir or lamivudine when used with other antiretrovirals.</p> <p><sup>2</sup>This adverse reaction was not identified to be reasonably attributed to Triumeq (dolutegravir + abacavir/lamivudine) in clinical trials, so the highest frequency category observed from the label of the individual components was used (e.g., for dolutegravir, abacavir and/or lamivudine).</p>	

#### Description of selected adverse reactions

##### *Hypersensitivity reactions*

Both abacavir and dolutegravir are associated with a risk for hypersensitivity reactions (HSR), which were observed more commonly with abacavir. Hypersensitivity reaction observed for each of these medicinal products (described below) share some common features such as fever and/or rash with other symptoms indicating multi-organ involvement. Time to onset was typically 10-14 days for both abacavir and dolutegravir-associated reactions, although reactions to abacavir may occur at any time during therapy. Treatment with Triumeq must be stopped without delay if HSR cannot be ruled out on clinical grounds, and therapy with Triumeq or other abacavir or dolutegravir containing products must never be re-initiated. Please refer to section 4.4 for further details on patient management in the event of a suspected HSR to Triumeq.

##### Dolutegravir hypersensitivity

Symptoms have included rash, constitutional findings, and sometimes, organ dysfunction, including severe liver reactions.

**Abacavir hypersensitivity**

The signs and symptoms of this HSR are listed below. These have been identified either from clinical studies or post marketing surveillance. Those reported in at least 10% of patients with a hypersensitivity reaction are in bold text.

Almost all patients developing hypersensitivity reactions will have fever and/or rash (usually maculopapular or urticarial) as part of the syndrome, however reactions have occurred without rash or fever. Other key symptoms include gastrointestinal, respiratory or constitutional symptoms such as lethargy and malaise.

<i>Skin</i>	<b>Rash</b> (usually maculopapular or urticarial)
<i>Gastrointestinal tract</i>	<b>Nausea, vomiting, diarrhoea, abdominal pain</b> , mouth ulceration
<i>Respiratory tract</i>	<b>Dyspnoea, cough</b> , sore throat, adult respiratory distress syndrome, respiratory failure
<i>Miscellaneous</i>	<b>Fever, lethargy, malaise</b> , oedema, lymphadenopathy, hypotension, conjunctivitis, anaphylaxis
<i>Neurological/Psychiatry</i>	<b>Headache</b> , paraesthesia
<i>Haematological</i>	Lymphopenia
<i>Liver/pancreas</i>	<b>Elevated liver function tests</b> , hepatitis, hepatic failure
<i>Musculoskeletal</i>	<b>Myalgia</b> , rarely myolysis, arthralgia, elevated creatine phosphokinase
<i>Urology</i>	Elevated creatinine, renal failure

Symptoms related to this HSR worsen with continued therapy and can be life-threatening and in rare instance, have been fatal.

Restarting abacavir following an abacavir HSR results in a prompt return of symptoms within hours. This recurrence of the HSR is usually more severe than on initial presentation, and may include life-threatening hypotension and death. Similar reactions have also occurred infrequently after restarting abacavir in patients who had only one of the key symptoms of hypersensitivity (see above) prior to stopping abacavir; and on very rare occasions have also been seen in patients who have restarted therapy with no preceding symptoms of a HSR (i.e., patients previously considered to be abacavir tolerant).

***Lactic acidosis***

Cases of lactic acidosis, sometimes fatal, usually associated with severe hepatomegaly and hepatic steatosis, have been reported with the use of nucleoside analogues (see section 4.4).

***Lipodystrophy***

Combination antiretroviral therapy (CART) has been associated with redistribution of body fat (lipodystrophy) in HIV patients including the loss of peripheral and facial subcutaneous fat, increased intra-abdominal and visceral fat, breast hypertrophy and dorsocervical fat accumulation (buffalo hump).

### *Metabolic abnormalities*

CART has been associated with metabolic abnormalities such as hypertriglyceridaemia, hypercholesterolaemia, insulin resistance, hyperglycaemia and hyperlactataemia (see section 4.4).

### *Osteonecrosis*

Cases of osteonecrosis have been reported, particularly in patients with generally acknowledged risk factors, advanced HIV disease or long-term exposure to CART. The frequency of this is unknown (see section 4.4).

### *Immune reactivation syndrome*

In HIV-infected patients with severe immune deficiency at the time of initiation of CART, an inflammatory reaction to asymptomatic or residual opportunistic infections may arise. Autoimmune disorders (such as Graves' disease) have also been reported; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment (see section 4.4).

### Changes in laboratory chemistries

Increases in serum creatinine occurred within the first week of treatment with dolutegravir and remained stable through 96 weeks. In the SINGLE study a mean change from baseline of 12.6 µmol/L was observed after 96 weeks of treatment. These changes are not considered to be clinically relevant since they do not reflect a change in glomerular filtration rate.

Asymptomatic creatine phosphokinase (CPK) elevations mainly in association with exercise have also been reported with dolutegravir therapy.

### Co-infection with Hepatitis B or C

In dolutegravir Phase III studies patients with hepatitis B and/or C co-infection were permitted to enrol provided that baseline liver chemistry tests did not exceed 5 times the upper limit of normal (ULN). Overall, the safety profile in patients co-infected with hepatitis B and/or C was similar to that observed in patients without hepatitis B or C co-infection, although the rates of AST and ALT abnormalities were higher in the subgroup with hepatitis B and/or C co-infection for all treatment groups.

### Paediatric population

There are no clinical study data on the effects of Triumeq in the paediatric population. Individual components have been investigated in adolescents (12 to 17 years).

Based on limited available data with the dolutegravir single entity used in combination with other antiretroviral agents to treat adolescents (12 to 17 years), there were no additional types of adverse reactions beyond those observed in the adult population.

The individual preparations of abacavir and lamivudine have been investigated separately, and as a dual nucleoside backbone, in combination antiretroviral therapy to treat ART- naive and ART-experienced HIV- infected paediatric patients (data available on the use of abacavir and lamivudine in infants less than three months are limited). No additional types of adverse reactions have been observed beyond those characterised for the adult population.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## 4.9 Overdose

No specific symptoms or signs have been identified following acute overdose with dolutegravir, abacavir or lamivudine, apart from those listed as adverse reactions.

Further management should be as clinically indicated or as recommended by the national poisons centre, where available. There is no specific treatment for an overdose of Triumeq. If overdose occurs, the patient should be treated supportively with appropriate monitoring, as necessary. Since lamivudine is dialysable, continuous haemodialysis could be used in the treatment of overdose, although this has not been studied. It is not known whether abacavir can be removed by peritoneal dialysis or haemodialysis. As dolutegravir is highly bound to plasma proteins, it is unlikely that it will be significantly removed by dialysis.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antivirals for systemic use, antivirals for treatment of HIV infections, combinations. ATC code: J05AR13

#### Mechanism of action

Dolutegravir inhibits HIV integrase by binding to the integrase active site and blocking the strand transfer step of retroviral Deoxyribonucleic acid (DNA) integration which is essential for the HIV replication cycle.

Abacavir and lamivudine are potent selective inhibitors of HIV-1 and HIV-2. Both abacavir and lamivudine are metabolised sequentially by intracellular kinases to the respective 5'-triphosphates (TP) which are the active moieties with extended intracellular half-lives supporting once daily dosing (see section 5.2). Lamivudine-TP (an analogue for cytidine) and carbovir-TP (the active triphosphate form of abacavir, an analogue for guanosine) are substrates for and competitive inhibitors of HIV reverse transcriptase (RT). However, their main antiviral activity is through incorporation of the monophosphate form into the viral DNA chain, resulting in chain termination. Abacavir and lamivudine triphosphates show significantly less affinity for host cell DNA polymerases.

#### Pharmacodynamic effects

##### *Antiviral activity in vitro*

Dolutegravir, abacavir and lamivudine have been shown to inhibit replication of lab-strains and clinical isolates of HIV in a number of cell types, including transformed T cell lines, monocyte/macrophage derived lines and primary cultures of activated peripheral blood mononuclear cells (PBMCs) and monocyte/macrophages. The concentration of drug necessary to effect viral replication by 50% (IC<sub>50</sub> - half maximal inhibitory concentration) varied according to virus and host cell type.

The IC<sub>50</sub> for dolutegravir in various lab-strains using PBMC was 0.5 nM, and when using MT-4 cells it ranged from 0.7-2 nM. Similar IC<sub>50</sub>s were seen for clinical isolates without any major difference

between subtypes; in a panel of 24 HIV-1 isolates of clades A, B, C, D, E, F and G and group O the mean IC<sub>50</sub> value was 0.2 nM (range 0.02-2.14). The mean IC<sub>50</sub> for 3 HIV-2 isolates was 0.18 nM (range 0.09-0.61).

The mean IC<sub>50</sub> for abacavir against lab-strains of HIV-1IIIB and HIV-1HXB2 ranged from 1.4 to 5.8 μM. The median or mean IC<sub>50</sub> values for lamivudine against lab-strains of HIV-1 ranged from 0.007 to 2.3 μM. The mean IC<sub>50</sub> against lab-strains of HIV-2 (LAV2 and EHO) ranged from 1.57 to 7.5 μM for abacavir and from 0.16 to 0.51 μM for lamivudine.

The IC<sub>50</sub> values of abacavir against HIV-1 Group M subtypes (A-G) ranged from 0.002 to 1.179 μM, against Group O from 0.022 to 1.21 μM, and against HIV-2 isolates, from 0.024 to 0.49 μM. For lamivudine, the IC<sub>50</sub> values against HIV-1 subtypes (A-G) ranged from 0.001 to 0.170 μM, against Group O from 0.030 to 0.160 μM and against HIV-2 isolates from 0.002 to 0.120 μM in peripheral blood mononuclear cells.

HIV-1 isolates (CRF01\_AE, n=12; CRF02\_AG, n=12; and Subtype C or CRF\_AC, n=13) from 37 untreated patients in Africa and Asia were susceptible to abacavir (IC<sub>50</sub> fold changes < 2.5), and lamivudine (IC<sub>50</sub> fold changes < 3.0), except for two CRF02\_AG isolates with fold changes of 2.9 and 3.4 for abacavir. Group O isolates from antiviral naïve patients tested for lamivudine activity were highly sensitive.

The combination of abacavir and lamivudine has demonstrated antiviral activity in cell culture against non-subtype B isolates and HIV-2 isolates with equivalent antiviral activity as for subtype B isolates.

#### *Antiviral activity in combination with other antiviral agents*

No antagonistic effects *in vitro* were seen with dolutegravir and other antiretrovirals (tested agents: stavudine, abacavir, efavirenz, nevirapine, lopinavir, amprenavir, enfuvirtide, maraviroc, adefovir and raltegravir). In addition, ribavirin had no apparent effect on dolutegravir activity.

The antiviral activity of abacavir in cell culture was not antagonized when combined with the nucleoside reverse transcriptase inhibitors (NRTIs) didanosine, emtricitabine, lamivudine, stavudine, tenofovir, zalcitabine or zidovudine, the non-nucleoside reverse transcriptase inhibitor (NNRTI) nevirapine, or the protease inhibitor (PI) amprenavir.

No antagonistic effects *in vitro* were seen with lamivudine and other antiretrovirals (tested agents: abacavir, didanosine, nevirapine, zalcitabine, and zidovudine).

#### *Effect of human serum*

In 100% human serum, the mean fold shift for dolutegravir activity was 75 fold, resulting in protein adjusted IC<sub>90</sub> of 0.064 ug/mL. Plasma protein binding studies *in vitro* indicate that abacavir binds only low to moderately (~49%) to human plasma proteins at therapeutic concentrations. Lamivudine exhibits linear pharmacokinetics over the therapeutic dose range and displays low plasma protein binding (less than 36%).

#### Resistance

##### *Resistance in vitro: (dolutegravir)*

Serial passage is used to study resistance evolution *in vitro*. When using the lab-strain HIVIII during passage over 112 days, mutations selected appeared slowly, with substitutions at positions S153Y and F. These mutations were not selected in patients treated with dolutegravir in the clinical studies.

Using strain NL432 mutations E92Q (fold change 3) and G193E (fold change 3) were selected. These mutations have been selected in patients with pre-existing raltegravir resistance and who were then treated with dolutegravir (listed as secondary mutations for dolutegravir).

In further selection experiments using clinical isolates of subtype B, mutation R263K was seen in all five isolates (after 20 weeks and onwards). In subtype C (n=2) and A/G (n=2) isolates the integrase substitution R263K was selected in one isolate, and G118R in two isolates. R263K was reported from two individual patients with subtype B and subtype C in the clinical program for ART experienced, INI naive subjects, but without effects on dolutegravir susceptibility *in vitro*. G118R lowers the susceptibility to dolutegravir in site directed mutants (fold change 10), but was not detected in patients receiving dolutegravir in the Phase III program.

Primary mutations for raltegravir/elvitegravir (Q148H/R/K, N155H, Y143R/H/C, E92Q, T66I) do not affect the *in vitro* susceptibility of dolutegravir as single mutations. When mutations listed as secondary integrase inhibitor associated mutations (for raltegravir/elvitegravir) are added to primary mutations (excluding at Q148) in experiments with site directed mutants, dolutegravir susceptibility remains at or near wildtype level. In the case of the Q148-mutation viruses, increasing dolutegravir fold change is seen as the number of secondary mutations increase. The effect of the Q148-based mutations (H/R/K) was also consistent with *in vitro* passage experiments with site directed mutants. In serial passage with strain NL432-based site directed mutants at N155H or E92Q, no further selection of resistance was seen (fold change unchanged around 1). In contrast, starting passage with mutants with mutation Q148H (fold change 1), a variety of raltegravir associated secondary mutations accumulated with a consequent increase of fold change to values >10. A clinically relevant phenotypic cut-off value (fold change vs wild type virus) has not been determined; genotypic resistance was a better predictor for outcome.

Seven hundred and five raltegravir resistant isolates from raltegravir experienced patients were analyzed for susceptibility to dolutegravir. Dolutegravir has a <10 fold change against 94% of the 705 clinical isolates.

#### *Resistance in vivo: (dolutegravir)*

In previously untreated patients receiving dolutegravir + 2 NRTIs in Phase IIb and Phase III, no development of resistance to the integrase class, or to the NRTI class was seen (n=876, follow-up of 48-96 weeks).

In patients with prior failed therapies, but naïve to the integrase class (SAILING study), integrase inhibitor substitutions were observed in 4/354 patients (follow-up 48 weeks) treated with dolutegravir, which was given in combination with an investigator selected background regimen (BR). Of these four, two subjects had a unique R263K integrase substitution, with a maximum fold change of 1.93, one subject had a polymorphic V151V/I integrase substitution, with maximum fold change of 0.92, and one subject had pre-existing integrase mutations and is assumed to have been integrase experienced or infected with integrase resistant virus by transmission. The R263K mutation was also selected *in vitro* (see above).

#### *Resistance in vitro and in vivo: (abacavir and lamivudine)*

Abacavir-resistant isolates of HIV-1 have been selected *in vitro* and *in vivo* and are associated with specific genotypic changes in the RT codon region (codons M184V, K65R, L74V and Y115F). During *in vitro* abacavir selection the M184V mutation occurred first and resulted in about a 2 fold increase in IC50, below the abacavir clinical cut-off of 4.5 fold change. Continued passage in increasing concentrations of drug resulted in selection for double RT mutants 65R/184V and 74V/184V or triple RT mutant 74V/115Y/184V. Two mutations conferred a 7 to 8 fold change in abacavir susceptibility and combinations of three mutations were required to confer more than an 8 fold change in susceptibility.

HIV-1 resistance to lamivudine involves the development of a M184I or M184V amino acid change close to the active site of the viral RT. This variant arises both *in vitro* and in HIV-1 infected patients

treated with lamivudine-containing antiretroviral therapy. M184V mutants display greatly reduced susceptibility to lamivudine and show diminished viral replicative capacity *in vitro*. M184V is associated with about a 2 fold increase in abacavir resistance but does not confer clinical resistance for abacavir.

Isolates resistant to abacavir may also show reduced sensitivity to lamivudine. The combination of abacavir/lamivudine has demonstrated decreased susceptibility to viruses with the substitutions K65R with or without the M184V/I substitution, and to viruses with L74V plus the M184V/I substitution.

Cross-resistance between dolutegravir or abacavir or lamivudine and antiretrovirals from other classes e.g. PIs or NNRTIs is unlikely.

#### Effects on electrocardiogram

No relevant effects were seen on the QTc interval, with doses of dolutegravir exceeding the clinical dose by approximately 3 fold. Similar studies were not conducted with either abacavir or lamivudine.

#### Clinical efficacy and safety

The efficacy of Triumeq in HIV-infected, therapy naive subjects is based on the analyses of data from two randomized, international, double-blind, active-controlled trials, SINGLE (ING114467) and SPRING-2 (ING113086) and the international, open-label, active-controlled trial FLAMINGO (ING114915).

In SINGLE, 833 patients were treated with dolutegravir 50 mg once daily plus fixed-dose abacavir-lamivudine (DTG + ABC/3TC) or fixed-dose efavirenz-tenofovir-emtricitabine (EFV/TDF/FTC). At baseline, median patient age was 35 years, 16% were female, 32% non-white, 7% had hepatitis C co-infection and 4% were CDC Class C, these characteristics were similar between treatment groups.

Week 48 outcomes (including outcomes by key baseline covariates) are shown in Table 3.

Table 3: Virologic Outcomes of Randomized Treatment of SINGLE at 48 Weeks (Snapshot algorithm)

	<b>48 weeks</b>	
	<b>DTG 50 mg + ABC/3TC once daily N=414</b>	<b>EFV/TDF/FTC once daily N=419</b>
<b>HIV-1 RNA &lt;50 copies/mL</b>	88%	81%
<b>Treatment Difference*</b>	7.4% (95% CI: 2.5%, 12.3%)	
<b>Virologic non response†</b>	5%	6%
<b>No virologic data at Weeks 48 window</b>	7%	13%
<b>Reasons</b>		
Discontinued study/study drug due to adverse event or death‡	2%	10%
Discontinued study/study drug for other reasons§	5%	3%
Missing data during window but on study	0	<1%
<b>HIV-1 RNA &lt;50 copies/mL by baseline covariates</b>		
<b>Baseline Plasma Viral Load (copies/mL)</b>	n / N (%)	n / N (%)
≤100,000	253 / 280 (90%)	238 / 288 (83%)
>100,000	111 / 134 (83%)	100 / 131 (76%)
<b>Baseline CD4+ (cells/ mm<sup>3</sup>)</b>		
<200	45 / 57 (79%)	48 / 62 (77%)
200 to <350	143 / 163 (88%)	126 / 159 (79%)
≥350	176 / 194 (91%)	164 / 198 (83%)
<b>Gender</b>		
Male	307 / 347 (88%)	291 / 356 (82%)
Female	57 / 67 (85%)	47 / 63 (75%)
<b>Race</b>		
White	255 / 284 (90%)	238 / 285 (84%)
African-American/African Heritage/Other	109 / 130 (84%)	99 / 133 (74%)
<b>Age (years)</b>		
<50	319 / 361 (88%)	302 / 375 (81%)
≥50	45 / 53 (85%)	36 / 44 (82%)

\* Adjusted for baseline stratification factors.

† Includes subjects who discontinued prior to Week 48 for lack or loss of efficacy and subjects who are ≥50 copies in the 48 week window.

‡ Includes subjects who discontinued due to an adverse event or death at any time point from Day 1 through the Week 48 analysis window if this resulted in no virologic data on treatment during the analysis window.

§ Includes reasons such as withdrew consent, loss to follow-up, moved, protocol deviation.

Notes: ABC/3TC = abacavir 600 mg, lamivudine 300 mg in the form of Kivexa/Epzicom fixed dose combination (FDC)

EFV/TDF/FTC = efavirenz 600 mg, tenofovir 300 mg, emtricitabine 200 mg in the form of Atripla FDC.

In the primary 48 weeks analysis, the proportion of patients with virologic suppression in the dolutegravir + ABC/3TC arm, was superior to the Efavirenz + TDF/FTC arm, p=0.003, the same treatment difference was observed in subjects defined by baseline HIV RNA level (< or > 100,000 copies/mL).

The median time to viral suppression was shorter with ABC/3TC + DTG (28 vs 84 days, p<0.0001). The adjusted mean change in CD4+ T cell count from baseline were 267 cells versus 208 cells/mm<sup>3</sup>, respectively (p<0.001). Both the time to viral suppression and change from baseline analyses were pre-specified and adjusted for multiplicity. At 96 weeks, the response was 80% vs 72%, respectively. The difference in the endpoint remained statistically significant (p=0.006). The statistically higher responses on DTG+ABC/3TC were driven by a higher rate of withdrawals due to AEs in the EFV/TDF/FTC arm, irrespective of viral load strata. Overall treatment differences at Week 96 are applicable to patients with high and low Baseline viral loads. At 144 weeks in the open-label phase of SINGLE, virologic suppression was maintained, the DTG +ABC/3TC arm (71%) was superior to the EFV/TDF/FTC arm (63%), treatment difference was 8.3% (2.0, 14.6).

In SPRING-2, 822 patients were treated with either dolutegravir 50 mg once daily or raltegravir 400 mg twice daily (blinded), both with fixed-dose ABC/3TC (around 40%) or TDF/FTC (around 60%), given open label. Baseline demographics and outcomes are summarised in Table 4. Dolutegravir was non-inferior to raltegravir, including within the subset of patients with the abacavir/lamivudine background regimen.

Table 4: Demographics and virologic outcomes of randomized treatment of SPRING-2 (snapshot algorithm)

	<b>DTG 50 mg once daily + 2 NRTI N=411</b>	<b>RAL 400mg twice daily + 2 NRTI N=411</b>
<b>Demographics</b>		
Median Age (years)	37	35
Female	15%	14%
Non-white	16%	14%
Hepatitis B and/or C	13%	11%
CDC class C	2%	2%
ABC/3TC backbone	41%	40%
<b>Week 48 efficacy results</b>		
HIV-1 RNA <50 copies/mL	88%	85%
Treatment difference*	2.5% (95% CI: -2.2%, 7.1%)	
Virologic non response†	5%	8%
No virologic data at Weeks 48 window	7%	7%
<b>Reasons</b>		
Discontinued study/study drug due to adverse event or death‡	2%	1%
Discontinued study/study drug for other reasons§	5%	6%
HIV-1 RNA <50 copies/mL for those on ABC/3TC	86%	87%
<b>Week 96 efficacy results</b>		
HIV-1 RNA <50 copies/mL	81%	76%
Treatment difference*	4.5% (95% CI: -1.1%, 10.0%)	
HIV-1 RNA <50 copies/mL for those on ABC/3TC	74%	76%
* Adjusted for baseline stratification factors.		
† Includes subjects who discontinued prior to Week 48 for lack or loss of efficacy and subjects who are ≥50 copies in the 48 week window.		
‡ Includes subjects who discontinued due to an adverse event or death at any time point from Day 1 through the Week 48 analysis window if this resulted in no virologic data on treatment during the analysis window.		
§ Includes reasons such as protocol deviation, lost to follow up, and withdrew consent.		
Notes: DTG = dolutegravir, RAL = raltegravir.		

In FLAMINGO, 485 patients were treated with dolutegravir 50 mg once daily or darunavir/ritonavir (DRV/r) 800 mg/100 mg once daily, both with ABC/3TC (around 33%) or TDF/FTC (around 67%). All treatments were given open-label. Main demographics and outcomes are summarised in Table 5.

Table 5: Demographics and Week 48 virologic outcomes of randomized treatment of FLAMINGO (snapshot algorithm)

	<b>DTG 50 mg once daily + 2 NRTI</b> <b>N=242</b>	<b>DRV+RTV 800mg + 100mg once daily +2 NRTI</b> <b>N=242</b>
<b>Demographics</b>		
Median Age (years)	34	34
Female	13%	17%
Non-white	28%	27%
Hepatitis B and/or C	11%	8%
CDC class C	4%	2%
ABC/3TC backbone	33%	33%
<b>Week 48 Efficacy Results</b>		
HIV-1 RNA <50 copies/mL	90%	83%
Treatment Difference*	7.1% (95% CI: 0.9%, 13.2%)	
Virologic non response†	6%	7%
No virologic data at Weeks 48 window	4%	10%
<u>Reasons</u>		
Discontinued study/study drug due to adverse event or death‡	1%	4%
Discontinued study/study drug for other reasons§	2%	5%
Missing data during window but on study	<1%	2%
HIV-1 RNA <50copies/mL for those on ABC/3TC	90%	85%
Median time to viral suppression**	28 days	85 days

\* Adjusted for baseline stratification factors, p=0.025.  
 † Includes subjects who discontinued prior to Week 48 for lack or loss of efficacy and subjects who are ≥50 copies in the 48 week window.  
 ‡ Includes subjects who discontinued due to an adverse event or death at any time point from Day 1 through the Week 48 analysis window if this resulted in no virologic data on treatment during the analysis window.  
 § Includes reasons such as withdrew consent, loss to follow-up, protocol deviation.  
 \*\* p<0.001.  
 Notes: DRV+RTV = darunavir + ritonavir, DTG = dolutegravir.

At 96 weeks, virologic suppression in the dolutegravir group (80%) was superior to the DRV/r group (68%), (adjusted treatment difference [DTG-(DRV+RTV)]: 12.4%; 95% CI: [4.7, 20.2]). Response rates at 96 weeks were 82% for DTG+ABC/3TC and 75% for DRV/r+ABC/3TC.

#### De novo resistance in patients failing therapy in SINGLE, SPRING-2 and FLAMINGO

*De novo* resistance was not detected to the integrase class or the NRTI class in any patients who were treated with dolutegravir + abacavir/lamivudine in the three studies mentioned.

For the comparators typical resistance was detected with TDF/FTC/EFV (SINGLE; six with NNRTI associated resistance and one with major NRTI resistance) and with 2 NRTIs + raltegravir (SPRING-2; four with major NRTI resistance and one with raltegravir resistance), while no *de novo* resistance was detected in patients treated with 2 NRTIs + DRV/RTV (FLAMINGO).

#### Paediatric population

In a Phase I/II 48 week multicentre, open-label study (P1093/ING112578), the pharmacokinetic parameters, safety, tolerability and efficacy of dolutegravir was evaluated in combination regimens in HIV-1 infected infants, children and adolescents.

At 24 weeks, 16 of 23 (69%) adolescents (12 to 17 years of age) treated with dolutegravir once daily (35 mg n=4; 50 mg n=19) plus OBR achieved viral load less than 50 copies/mL.

Twenty out of 23 children and adolescents (87%) had  $>1 \log_{10}$  c/mL decrease from Baseline in HIV-1 RNA or HIV-1 RNA <400 c/mL at Week 24. Four subjects had virologic failure none of which had INI resistance at the time of virologic failure.

## 5.2 Pharmacokinetic properties

The Triumeq tablet has been shown to be bioequivalent to dolutegravir single entity tablet and abacavir/lamivudine fixed-dose combination tablet (ABC/3TC FDC) administered separately. This was demonstrated in a single dose, 2-way crossover bioequivalence study of Triumeq (fasted) versus 1 x 50 mg dolutegravir tablet, plus 1 x 600mg abacavir/300 mg lamivudine tablet (fasted) in healthy subjects (n=66). The effect of a high fat meal on the Triumeq tablet was evaluated in a subgroup of subjects in this study (n=12). Plasma  $C_{max}$  and AUC of dolutegravir following administration of Triumeq with a high fat meal were 37% and 48% higher, respectively, than those following administration of Triumeq in the fasted state. This is not considered clinically significant (see Absorption). The effect of food on plasma exposures of abacavir and lamivudine following administration of Triumeq with a high fat meal were very similar to prior food effects observed with ABC/3TC FDC. These results indicate that Triumeq can be taken with or without food.

The pharmacokinetic properties of dolutegravir, lamivudine and abacavir are described below.

### Absorption

Dolutegravir, abacavir and lamivudine are rapidly absorbed following oral administration. The absolute bioavailability of dolutegravir has not been established. The absolute bioavailability of oral abacavir and lamivudine in adults is about 83% and 80-85% respectively. The mean time to maximal serum concentrations ( $t_{max}$ ) is about 2 to 3 hours (post dose for tablet formulation), 1.5 hours and 1.0 hour for dolutegravir, abacavir and lamivudine, respectively.

Exposure to dolutegravir was generally similar between healthy subjects and HIV-1-infected subjects. In HIV-1-infected adult subjects following dolutegravir 50 mg once daily, the steady-state pharmacokinetic parameters (geometric mean [%CV]) based on population pharmacokinetic analyses were  $AUC_{(0-24)} = 53.6$  (27)  $\mu\text{g.h}/\text{mL}$ ,  $C_{max} = 3.67$  (20)  $\mu\text{g}/\text{mL}$ , and  $C_{min} = 1.11$  (46)  $\mu\text{g}/\text{mL}$ . Following a single dose of 600 mg of abacavir, the mean (CV)  $C_{max}$  is 4.26  $\mu\text{g}/\text{ml}$  (28%) and the mean (CV)  $AUC_{\infty}$  is 11.95  $\mu\text{g.h}/\text{ml}$  (21%). Following multiple-dose oral administration of lamivudine 300 mg once daily for seven days, the mean (CV) steady-state  $C_{max}$  is 2.04  $\mu\text{g}/\text{ml}$  (26%) and the mean (CV)  $AUC_{24}$  is 8.87  $\mu\text{g.h}/\text{ml}$  (21%).

Plasma  $C_{max}$  and AUC of dolutegravir following administration of Triumeq with a high fat meal were 37% and 48% higher, respectively, than those following administration of Triumeq in the fasted state). For abacavir there was a decrease in  $C_{max}$  with 23% and AUC was unchanged. The exposure of lamivudine was similar with and without food. These results indicate that Triumeq can be taken with or without food.

## Distribution

The apparent volume of distribution of dolutegravir (following oral administration of suspension formulation, Vd/F) is estimated at 12.5 L. Intravenous studies with abacavir and lamivudine showed that the mean apparent volume of distribution is 0.8 and 1.3 l/kg respectively.

Dolutegravir is highly bound (> 99%) to human plasma proteins based on *in vitro* data. Binding of dolutegravir to plasma proteins is independent of dolutegravir concentration. Total blood and plasma drug-related radioactivity concentration ratios averaged between 0.441 to 0.535, indicating minimal association of radioactivity with blood cellular components. The unbound fraction of dolutegravir in plasma is increased at low levels of serum albumin (<35 g/L) as seen in subjects with moderate hepatic impairment. Plasma protein binding studies *in vitro* indicate that abacavir binds only low to moderately (~49%) to human plasma proteins at therapeutic concentrations. Lamivudine exhibits linear pharmacokinetics over the therapeutic dose range and displays limited plasma protein binding *in vitro* (< 36%).

Dolutegravir, abacavir and lamivudine are present in cerebrospinal fluid (CSF).

In 13 treatment-naïve subjects on a stable dolutegravir plus abacavir/lamivudine regimen, dolutegravir concentration in CSF averaged 18 ng/mL (comparable to unbound plasma concentration, and above the IC50). Studies with abacavir demonstrate a CSF to plasma AUC ratio of between 30 to 44%. The observed values of the peak concentrations are 9 fold greater than the IC50 of abacavir of 0.08 µg/ml or 0.26 µM when abacavir is given at 600 mg twice daily. The mean ratio of CSF/serum lamivudine concentrations 2-4 hours after oral administration was approximately 12%. The true extent of CNS penetration of lamivudine and its relationship with any clinical efficacy is unknown.

Dolutegravir is present in the female and male genital tract. AUC in cervicovaginal fluid, cervical tissue and vaginal tissue were 6-10% of those in corresponding plasma at steady state. AUC in semen was 7% and 17% in rectal tissue of those in corresponding plasma at steady state.

## Biotransformation

Dolutegravir is primarily metabolized via UGT1A1 with a minor CYP3A component (9.7% of total dose administered in a human mass balance study). Dolutegravir is the predominant circulating compound in plasma; renal elimination of unchanged active substance is low (< 1% of the dose). Fifty-three percent of total oral dose is excreted unchanged in the faeces. It is unknown if all or part of this is due to unabsorbed active substance or biliary excretion of the glucuronide conjugate, which can be further degraded to form the parent compound in the gut lumen. Thirty-two percent of the total oral dose is excreted in the urine, represented by ether glucuronide of dolutegravir (18.9% of total dose), N-dealkylation metabolite (3.6% of total dose), and a metabolite formed by oxidation at the benzylic carbon (3.0% of total dose).

Abacavir is primarily metabolised by the liver with approximately 2% of the administered dose being renally excreted, as unchanged compound. The primary pathways of metabolism in man are by alcohol dehydrogenase and by glucuronidation to produce the 5'-carboxylic acid and 5'-glucuronide which account for about 66% of the administered dose. These metabolites are excreted in the urine.

Metabolism of lamivudine is a minor route of elimination. Lamivudine is predominately cleared by renal excretion of unchanged lamivudine. The likelihood of metabolic drug interactions with lamivudine is low due to the small extent of hepatic metabolism (5-10%).

## Drug interactions

*In vitro*, dolutegravir demonstrated no direct, or weak inhibition ( $IC_{50} > 50 \mu M$ ) of the enzymes cytochrome P<sub>450</sub> (CYP)1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6 CYP3A, uridine diphosphate glucuronosyl transferase (UGT)1A1 or UGT2B7, or the transporters PgP, BCRP, BSEP, OATP1B1, OATP1B3, OCT1, MATE2-K, MRP2 or MRP4. *In vitro*, dolutegravir did not induce CYP1A2, CYP2B6 or CYP3A4. Based on this data, dolutegravir is not expected to affect the pharmacokinetics of medicinal products that are substrates of major enzymes or transporters (see section 4.5).

*In vitro*, dolutegravir was not a substrate of human OATP 1B1, OATP 1B3 or OCT 1.

#### Elimination

Dolutegravir has a terminal half-life of ~14 hours. The apparent oral clearance (CL/F) is approximately 1 L/hr in HIV-infected patients based on a population pharmacokinetic analysis.

The mean half-life of abacavir is about 1.5 hours. The geometric mean terminal half-life of intracellular active moiety carbovirtriphosphate (TP) at steady-state is 20.6 hours. Following multiple oral doses of abacavir 300 mg twice a day there is no significant accumulation of abacavir. Elimination of abacavir is via hepatic metabolism with subsequent excretion of metabolites primarily in the urine. The metabolites and unchanged abacavir account for about 83% of the administered abacavir dose in the urine. The remainder is eliminated in the faeces.

The observed lamivudine half-life of elimination is 5 to 7 hours. For patients receiving lamivudine 300 mg once daily, the terminal intracellular half-life of lamivudine-TP was 16 to 19 hours. The mean systemic clearance of lamivudine is approximately 0.32 l/h/kg, predominantly by renal clearance (> 70%) via the organic cationic transport system. Studies in patients with renal impairment show lamivudine elimination is affected by renal dysfunction. Dose reduction is required for patients with creatinine clearance < 50 ml/min (see section 4.2).

#### Pharmacokinetic/pharmacodynamic relationship(s)

In a randomized, dose-ranging trial, HIV-1-infected subjects treated with dolutegravir monotherapy (ING111521) demonstrated rapid and dose-dependent antiviral activity with mean decline in HIV-1 RNA of  $2.5 \log_{10}$  at day 11 for 50 mg dose. This antiviral response was maintained for 3 to 4 days after the last dose in the 50 mg group.

#### Intracellular pharmacokinetics

The geometric mean terminal carbovir-TP intracellular half-life at steady-state was 20.6 hours, compared to the geometric mean abacavir plasma half-life of 2.6 hours. The terminal intracellular half-life of lamivudine-TP was prolonged to 16-19 hours, compared to the plasma lamivudine half-life of 5-7 hours, supporting once daily dosing of ABC and 3TC.

#### Special patient populations

##### *Hepatic impairment*

Pharmacokinetic data has been obtained for dolutegravir, abacavir and lamivudine separately.

Dolutegravir is primarily metabolized and eliminated by the liver. A single dose of 50 mg of dolutegravir was administered to 8 subjects with moderate hepatic impairment (Child-Pugh class B) and to 8 matched healthy adult controls. While the total dolutegravir concentration in plasma was similar, a 1.5 to 2 fold increase in unbound exposure to dolutegravir was observed in subjects with moderate hepatic impairment compared to healthy controls. No dosage adjustment is considered

necessary for patients with mild to moderate hepatic impairment. The effect of severe hepatic impairment on the pharmacokinetics of dolutegravir has not been studied.

Abacavir is metabolised primarily by the liver. The pharmacokinetics of abacavir have been studied in patients with mild hepatic impairment (Child-Pugh score 5-6) receiving a single 600 mg dose. The results showed that there was a mean increase of 1.89 fold [1.32; 2.70] in the abacavir AUC, and 1.58 [1.22; 2.04] fold in the elimination half-life. No recommendation on dose reduction is possible in patients with mild hepatic impairment due to substantial variability of abacavir exposure.

Data obtained in patients with moderate to severe hepatic impairment show that lamivudine pharmacokinetics are not significantly affected by hepatic dysfunction.

Based on data obtained for abacavir, Triumeq is not recommended in patients with moderate and severe hepatic impairment.

#### *Renal impairment*

Pharmacokinetic data have been obtained for dolutegravir, lamivudine and abacavir separately.

Renal clearance of unchanged active substance is a minor pathway of elimination for dolutegravir. A study of the pharmacokinetics of dolutegravir was performed in subjects with severe renal impairment ( $\text{CL}_{\text{cr}} < 30 \text{ mL/min}$ ). No clinically important pharmacokinetic differences between subjects with severe renal impairment ( $\text{CL}_{\text{cr}} < 30 \text{ mL/min}$ ) and matching healthy subjects were observed. Dolutegravir has not been studied in patients on dialysis, though differences in exposure are not expected.

Abacavir is primarily metabolised by the liver with approximately 2% of abacavir excreted unchanged in the urine. The pharmacokinetics of abacavir in patients with end-stage renal disease is similar to patients with normal renal function.

Studies with lamivudine show that plasma concentrations (AUC) are increased in patients with renal dysfunction due to decreased clearance.

Based on the lamivudine data, Triumeq is not recommended for patients with creatinine clearance of  $< 50 \text{ ml/min}$ .

#### Elderly

Population pharmacokinetic analysis of dolutegravir using data in HIV-1 infected adults showed that there was no clinically relevant effect of age on dolutegravir exposure.

Pharmacokinetic data for dolutegravir, abacavir and lamivudine in subjects  $> 65$  years of age are limited.

#### Paediatric population

The pharmacokinetics of dolutegravir in 10 antiretroviral treatment-experienced HIV-1 infected adolescents (12 to 17 years) showed that dolutegravir 50 mg once daily dosage resulted in dolutegravir exposure comparable to that observed in adults who received dolutegravir 50 mg once daily.

Limited data are available in adolescents receiving a daily dose of 600 mg of abacavir and 300 mg of lamivudine. Pharmacokinetic parameters are comparable to those reported in adults.

### Polymorphisms in drug metabolising enzymes

There is no evidence that common polymorphisms in drug metabolising enzymes alter dolutegravir pharmacokinetics to a clinically meaningful extent. In a meta-analysis using pharmacogenomics samples collected in clinical studies in healthy subjects, subjects with UGT1A1 (n=7) genotypes conferring poor dolutegravir metabolism had a 32% lower clearance of dolutegravir and 46% higher AUC compared with subjects with genotypes associated with normal metabolism via UGT1A1 (n=41).

### Gender

Population PK analyses using pooled pharmacokinetic data from Phase IIb and Phase III adult trials revealed no clinically relevant effect of gender on the exposure of dolutegravir. There is no evidence that a dose adjustment of dolutegravir, abacavir or lamivudine would be required based on the effects of gender on PK parameters.

### Race

Population PK analyses using pooled pharmacokinetic data from Phase IIb and Phase III adult trials revealed no clinically relevant effect of race on the exposure of dolutegravir. The pharmacokinetics of dolutegravir following single dose oral administration to Japanese subjects appear similar to observed parameters in Western (US) subjects. There is no evidence that a dose adjustment of dolutegravir, abacavir or lamivudine would be required based on the effects of race on PK parameters.

### Co-infection with Hepatitis B or C

Population pharmacokinetic analysis indicated that hepatitis C virus co-infection had no clinically relevant effect on the exposure to dolutegravir. There are limited pharmacokinetic data on subjects with hepatitis B co-infection (see section 4.4).

## **5.3 Preclinical safety data**

There are no data available on the effects of the combination of dolutegravir, abacavir and lamivudine in animals, except a negative *in vivo* rat micronucleus test which tested the effects of the combination of abacavir and lamivudine.

### Mutagenicity and carcinogenicity

Dolutegravir was not mutagenic or clastogenic using *in vitro* tests in bacteria and cultured mammalian cells, and an *in vivo* rodent micronucleus assay.

Neither abacavir nor lamivudine were mutagenic in bacterial tests, but consistent with other nucleoside analogues, inhibit cellular DNA replication in *in vitro* mammalian tests such as the mouse lymphoma assay. The results of an *in vivo* rat micronucleus test with abacavir and lamivudine in combination were negative.

Lamivudine has not shown any genotoxic activity in the *in vivo* studies. Abacavir has a weak potential to cause chromosomal damage both *in vitro* and *in vivo* at high tested concentrations.

The carcinogenic potential of a combination of dolutegravir, abacavir and lamivudine has not been tested. Dolutegravir was not carcinogenic in long term studies in the mouse and rat. In long-term oral carcinogenicity studies in rats and mice, lamivudine did not show any carcinogenic potential. Carcinogenicity studies with orally administered abacavir in mice and rats showed an increase in the

incidence of malignant and non-malignant tumours. Malignant tumours occurred in the preputial gland of males and the clitoral gland of females of both species, and in rats in the thyroid gland of males and in the liver, urinary bladder, lymph nodes and the subcutis of females.

The majority of these tumours occurred at the highest abacavir dose of 330 mg/kg/day in mice and 600 mg/kg/day in rats. The exception was the preputial gland tumour which occurred at a dose of 110 mg/kg in mice. The systemic exposure at the no effect level in mice and rats was equivalent to 3 and 7 times the human systemic exposure during therapy. While the clinical relevance of these findings is unknown, these data suggest that a potential carcinogenic risk to humans is outweighed by the clinical benefit.

#### Repeat-dose toxicity

The effect of prolonged daily treatment with high doses of dolutegravir has been evaluated in repeat oral dose toxicity studies in rats (up to 26 weeks) and in monkeys (up to 38 weeks). The primary effect of dolutegravir was gastrointestinal intolerance or irritation in rats and monkeys at doses that produce systemic exposures approximately 38 and 1.5 times the 50 mg human clinical exposure based on AUC, respectively. Because gastrointestinal (GI) intolerance is considered to be due to local active substance administration, mg/kg or mg/m<sup>2</sup> metrics are appropriate determinates of safety cover for this toxicity. GI intolerance in monkeys occurred at 30 times the human mg/kg equivalent dose (based on 50 kg human), and 11 times the human mg/m<sup>2</sup> equivalent dose for a total daily clinical dose of 50 mg.

In toxicology studies abacavir was shown to increase liver weights in rats and monkeys. The clinical relevance of this is unknown. There is no evidence from clinical studies that abacavir is hepatotoxic. Additionally, autoinduction of abacavir metabolism or induction of the metabolism of other medicinal products hepatically metabolised has not been observed in humans.

Mild myocardial degeneration in the heart of mice and rats was observed following administration of abacavir for two years. The systemic exposures were equivalent to 7 to 21 times the expected systemic exposure in humans. The clinical relevance of this finding has not been determined.

#### Reproductive toxicology

In reproductive toxicity studies in animals, dolutegravir, lamivudine and abacavir were shown to cross the placenta.

Oral administration of dolutegravir to pregnant rats at doses up to 1000 mg/kg daily from days 6 to 17 of gestation did not elicit maternal toxicity, developmental toxicity or teratogenicity (50 times the 50 mg human clinical exposure when administered in combination with abacavir and lamivudine based on AUC).

Oral administration of dolutegravir to pregnant rabbits at doses up to 1000 mg/kg daily from days 6 to 18 of gestation did not elicit developmental toxicity or teratogenicity (0.74 times the 50mg human clinical exposure when administered in combination with abacavir and lamivudine based on AUC). In rabbits, maternal toxicity (decreased food consumption, scant/no faeces/urine, suppressed body weight gain) was observed at 1000 mg/kg (0.74 times the 50 mg human clinical exposure when administered in combination with abacavir and lamivudine based on AUC).

Lamivudine was not teratogenic in animal studies but there were indications of an increase in early embryonic deaths in rabbits at relatively low systemic exposures, comparable to those achieved in humans. A similar effect was not seen in rats even at very high systemic exposure.

Abacavir demonstrated toxicity to the developing embryo and foetus in rats, but not in rabbits. These findings included decreased foetal body weight, foetal oedema, and an increase in skeletal variations/malformations, early intra-uterine deaths and still births. No conclusion can be drawn with regard to the teratogenic potential of abacavir because of this embryo-foetal toxicity.

Fertility studies in rats have shown that dolutegravir, abacavir and lamivudine have no effect on male or female fertility.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Mannitol (E421)  
Microcrystalline cellulose  
Povidone K29/32  
Sodium starch glycollate  
Magnesium stearate

#### Tablet coating

Opadry II Purple 85F90057 containing:  
Polyvinyl alcohol – part hydrolyzed  
titanium dioxide  
macrogol  
talc  
iron oxide black  
iron oxide red

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

2 years

### **6.4 Special precautions for storage**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

This medicinal product does not require any special temperature storage conditions.

### **6.5 Nature and contents of container**

White HDPE (high density polyethylene) bottles closed with polypropylene child-resistant closures, with a polyethylene faced induction heat seal liner. Each bottle contains 30 film-coated tablets and a desiccant.

Multipacks containing 90 (3 packs of 30) film-coated tablets. Each pack of 30 film-coated tablets contains a desiccant.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for disposal**

No special requirements.

### **7. MARKETING AUTHORISATION HOLDER**

ViiV Healthcare UK Limited  
980 Great West Road  
Brentford  
Middlesex  
TW8 9GS  
United Kingdom

### **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/940/001  
EU/1/14/940/002

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorization: 2<sup>nd</sup> September 2014

### **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

## **ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer(s) responsible for batch release

GLAXO WELLCOME, S.A.  
Avda. Extremadura, 3  
Pol. Ind. Allendeduero  
Aranda de Duero  
Burgos 09400  
Spain

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports**

The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation. Subsequently, the marketing authorisation holder shall submit periodic safety update reports for this product in accordance with the requirements set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

If the submission of a PSUR and the update of a RMP coincide, they can be submitted at the same time.

- **Additional risk minimisation measures**

The Marketing Authorisation Holder (MAH) shall ensure that all physicians who are expected to prescribe Triumeq are provided with a health care professional information pack containing the following:

- The Summary of Product Characteristics
- ABC HSR educational material for health care professionals

**Key elements included in the educational material** to increase understanding and awareness of ABC HSR and expand on the information already included in the SmPC:

**1. Major symptoms associated with ABC HSR** are fever (~80%), rash (~70%), gastrointestinal symptoms (>50%) such as nausea, abdominal pain, vomiting, and diarrhoea, generalise malaise, fatigue, and headache (~50%) and other symptoms (~30%) such as respiratory, mucosal, and musculoskeletal symptoms.

**Based on the above patients are advised to contact their physician immediately to determine whether they should stop taking abacavir if:**

- presence of skin rash; OR
- development of 1 or more symptom from at least 2 of the following groups:
- Fever
- Shortness of breath, sore throat or cough
- Nausea or vomiting or diarrhoea or abdominal pain
- Extreme tiredness or achiness or generally ill feeling

## **2. Risk factors for ABC HSR**

HLA-B\*5701 is the only identified pharmacogenetic marker that is consistently associated with clinical diagnosis of an ABC HSR reaction. However, some patients with a suspected ABC hypersensitivity reaction may not have the HLA-B\*5701 allele.

## **3. Recommendations for HLA-B\*5701 screening**

Before initiating abacavir therapy, clinicians should screen for HLA-B\*5701(in settings where validated screening methods are available). Clinical diagnosis of suspected hypersensitivity to ABC remains the basis for clinical decision making. HLA-B\*5701 screening for risk of ABC hypersensitivity should never be substituted for appropriate clinical vigilance and patient management in individuals receiving ABC. If ABC hypersensitivity cannot be ruled out on clinical grounds, ABC should be permanently discontinued and should not be restarted, regardless of the results of HLA-B\*5701 screening. Screening is also recommended prior to re-initiation of abacavir in patients of unknown HLA-B\*5701 status who have previously tolerated abacavir.

## **4. Information on HLA-B\*5701 testing**

The one-time HLA-B\*5701 test identifies people at high risk for this serious allergic reaction. The gold standards for HLA-B\*5701 screening are sequence-based genotyping and polymerase chain reaction sequencing of specific oligonucleotide probes. Blood or saliva samples are collected and tested for genetic sequences coding for the HLA-B\*5701 allele. Results of PREDICT-1 and SHAPE studies show that the presence of the HLA-B\*5701 allele is associated with increased risk of ABC hypersensitivity, regardless of race, screening for HLA-B\*5701 before starting treatment with ABC may identify subjects at increased risk of a HSR, avoiding treatment with ABC in subjects with the

HLA-B\*5701 allele was shown to significantly reduce the incidence rate of clinically diagnosed cases of hypersensitivity. Data from these studies do not support the use of skin patch testing in routine clinical practice. Only patients found to lack the HLA-B\*5701 allele should begin therapy with ABC.

## **5. Management of ABC HSR reaction**

Symptoms can occur at any time during treatment with ABC, but usually occur within the first 6 weeks of therapy. Symptoms are initially mild and evolve over days, becoming more severe with continued ABC therapy. Symptoms improve on cessation of ABC. Rechallenge can result in a more rapid and severe reaction, which can be fatal, therefore rechallenge is contraindicated.

## **6. Hypersensitivity case studies**

The educational material includes 3 case studies to demonstrate different clinical scenarios and their management.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****BOTTLE CARTON (INDIVIDUAL PACKS ONLY)****1. NAME OF THE MEDICINAL PRODUCT**

Triumeq 50 mg/600 mg/300 mg film-coated tablets  
dolutegravir/abacavir/lamivudine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains  
50 mg dolutegravir (as sodium),  
600 mg abacavir (as sulfate),  
300 mg lamivudine.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

30 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Detach enclosed Alert Card, it contains important safety information.

**WARNING**

In case of any symptoms suggesting hypersensitivity reactions, contact your doctor IMMEDIATELY.

Push here (with Alert card attached)

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

ViiV Healthcare UK Limited  
980 Great West Road  
Brentford  
Middlesex  
TW8 9GS  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/940/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

trumeq

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****BOTTLE CARTON (MULITPACKS ONLY – WITH BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Triumeq 50 mg/600 mg/300 mg film-coated tablets  
dolutegravir/abacavir/lamivudine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains  
50 mg dolutegravir (as sodium),  
600 mg abacavir (as sulfate),  
300 mg lamivudine.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

Multipack: 90 (3 packs of 30) film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

WARNING! In case of any symptoms suggesting hypersensitivity reactions, contact your doctor IMMEDIATELY.

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

ViiV Healthcare UK Limited  
980 Great West Road  
Brentford  
Middlesex  
TW8 9GS  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/940/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

triumeq

**PARTICULARS TO APPEAR ON THE INTERMEDIATE PACKAGING****INTERMEDIATE CARTON (WITHOUT BLUE BOX – COMPONENT OF MULTIPACK)****1. NAME OF THE MEDICINAL PRODUCT**

Triumeq 50 mg/600 mg/300 mg film-coated tablets  
dolutegravir/abacavir/lamivudine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains:  
50 mg dolutegravir (as sodium),  
600 mg abacavir (as sulfate),  
300 mg lamivudine.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

30 film-coated tablets. Component of a multipack, not to be sold separately.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Detach enclosed Alert Card, it contains important safety information.

**WARNING**

In case of any symptoms suggesting hypersensitivity reactions, contact your doctor IMMEDIATELY.

Push here (with Alert card attached)

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

ViiV Healthcare UK Limited  
980 Great West Road  
Brentford  
Middlesex  
TW8 9GS  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/940/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

triumeq

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING****BOTTLE LABEL****1. NAME OF THE MEDICINAL PRODUCT**

Triumeq 50 mg/600 mg/300 mg tablets  
dolutegravir/abacavir/lamivudine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains  
50 mg dolutegravir (as sodium),  
600 mg abacavir (as sulfate),  
300 mg lamivudine.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

30 tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

ViiV Healthcare UK Limited

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/940/001  
EU/1/14/940/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

## **TRIUMEQ TABLETS PATIENT ALERT CARD**

### **SIDE 1**

**IMPORTANT - PATIENT ALERT CARD**  
**Triumeq (dolutegravir / abacavir / lamivudine) tablets**  
**Carry this card with you at all times**

Since Triumeq contains abacavir some patients taking Triumeq may develop a hypersensitivity reaction (serious allergic reaction) which **can be life-threatening** if treatment with Triumeq is continued. **CONTACT YOUR DOCTOR IMMEDIATELY** for advice on whether you should stop taking Triumeq if:

- 1) you get a skin rash OR**
- 2) you get one or more symptoms from at least TWO of the following groups**
  - fever
  - shortness of breath, sore throat or cough
  - nausea or vomiting or diarrhoea or abdominal pain
  - severe tiredness or achiness or generally feeling ill

If you have discontinued Triumeq due to this reaction, **YOU MUST NEVER TAKE** Triumeq, or any medicine containing abacavir again as **within hours** you may experience a life-threatening lowering of your blood pressure or death.

(see reverse of card)

### **SIDE 2**

You should immediately contact your doctor if you think you are having a hypersensitivity reaction to Triumeq. Write your doctor's details below:

Doctor:..... Tel:.....

**If your doctor is not available, you must urgently seek alternative medical advice (e.g. the emergency unit of the nearest hospital).**

For general Triumeq information enquiries, contact:

**B. PACKAGE LEAFLET**

## **Package leaflet: Information for the patient**

### **Triumeq 50 mg/600 mg/300 mg film-coated tablets dolutegravir/abacavir/lamivudine**

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

#### **Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Triumeq is and what it is used for
2. What you need to know before you take Triumeq
3. How to take Triumeq
4. Possible side effects
5. How to store Triumeq
6. Contents of the pack and other information

#### **1. What Triumeq is and what it is used for**

Triumeq is a medicine that contains three active ingredients used to treat HIV infection: abacavir, lamivudine and dolutegravir. Abacavir and lamivudine belong to a group of anti-retroviral medicines called *nucleoside analogue reverse transcriptase inhibitors (NRTIs)*, and dolutegravir belongs to a group of anti-retroviral medicines called *integrase inhibitors (INIs)*.

Triumeq is used to treat **HIV (human immunodeficiency virus) infection** in adults and children over 12 years old who weigh at least 40 kg.

Before you are prescribed Triumeq your doctor will arrange a test to find out whether you carry a particular type of gene called HLA-B\*5701. Triumeq should not be used in patients who are known to carry the HLA-B\*5701 gene. Patients with this gene are at a high risk of developing a serious hypersensitivity (allergic) reaction if they use Triumeq (see 'hypersensitivity reactions' in section 4).

Triumeq does not cure HIV infection; it reduces the amount of virus in your body, and keeps it at a low level. It also increases the number of CD4 cells in your blood. CD4 cells are a type of white blood cells that are important in helping your body to fight infection.

Not everyone responds to treatment with Triumeq in the same way. Your doctor will monitor the effectiveness of your treatment.

#### **2. What you need to know before you take Triumeq**

##### **Do not take Triumeq:**

- if you are **allergic** (*hypersensitive*) to dolutegravir, abacavir (or any other medicine containing abacavir), or lamivudine, or any of the other ingredients of this medicine (listed in section 6).  
**Carefully read all the information about hypersensitivity reactions in Section 4.**
- if you are taking a medicine called **dofetilide** (to treat heart conditions).  
→ If you think any of these apply to you, tell your doctor.

## **Warnings and Precautions**

### **IMPORTANT — Hypersensitivity reactions**

**Triumeq contains abacavir and dolutegravir.** Both of these active ingredients can cause a serious allergic reaction known as a hypersensitivity reaction, which can be life threatening in people who continue to take abacavir-containing products.

**You must carefully read all the information under ‘Hypersensitivity reactions’ in the panel in Section 4.**

The Triumeq pack includes an **Alert Card** to remind you and medical staff about hypersensitivity. **Detach this card and keep it with you at all times.**

Some people taking Triumeq or other combination treatments for HIV are more at risk of serious side effects than others. You need to be aware of the extra risks:

- if you have ever had **liver disease**, including hepatitis B or C (if you have hepatitis B infection, don't stop Triumeq without your doctor's advice, as your hepatitis may come back)
- if you have a kidney problem  
→ **Talk to your doctor if any of these apply to you before using Triumeq.** You may need extra check-ups, including blood tests, while you're taking your medicine. See Section 4 for more information.

### Abacavir hypersensitivity reactions

Even patients who don't have the HLA-B\*5701 gene may still develop a **hypersensitivity reaction** (a serious allergic reaction).

→ **Carefully read all the information about hypersensitivity reactions in Section 4 of this leaflet.**

### Risk of heart attack

The possibility that abacavir increases the risk of having a heart attack cannot be excluded.

→ **Tell your doctor** if you have heart problems, if you smoke, or have other illnesses that may increase your risk of heart disease such as high blood pressure or diabetes. Don't stop taking Triumeq unless your doctor advises you to do so.

### Look out for important symptoms

Some people taking medicines for HIV infection develop other conditions, which can be serious. These include:

- symptoms of infections and inflammation
- joint pain, stiffness and bone problems

You need to know about important signs and symptoms to look out for while you're taking Triumeq.

→ **Read the information ‘Other possible side effects of combination therapy for HIV’ in Section 4 of this leaflet.**

### Protect other people

HIV infection is spread by sexual contact with someone who has the infection, or by transfer of infected blood (for example, by sharing injection needles). You can still pass on HIV when taking this medicine, although the risk is lowered by effective antiretroviral therapy. Discuss with your doctor the precautions needed to avoid infecting other people.

## **Children**

This medicine is not for children under 12 years of age. The use of Triumeq in children under 12 years of age has not yet been studied.

## **Other medicines and Triumeq**

Tell your doctor if you are taking, have recently taken or might take any other medicines, including herbal medicines and other medicines bought without a prescription.

Don't take Triumeq with the following medicine:

- dofetilide, used to treat **heart conditions**

Some medicines can affect how Triumeq works, or make it more likely that you will have side effects. Triumeq can also affect how some other medicines work.

**Tell your doctor** if you are taking any of the medicines *in the following list*:

- metformin, to treat **diabetes**
- medicines called **antacids**, to treat **indigestion** and **heartburn**. **Do not take an antacid** during the 6 hours before you take Triumeq, or for at least 2 hours after you take it. (*See also Section 3*).
- calcium supplements, iron supplements and multivitamins. **Do not take a calcium supplement, iron supplement or multivitamin** during the 6 hours before you take Triumeq, or for at least 2 hours after you take it (*see also Section 3*).
- emtricitabine, etravirine, efavirenz, nevirapine or tipranavir/ritonavir, to treat **HIV infection**
- other medicines containing lamivudine, used to treat **HIV infection** or **hepatitis B infection**
- cladribine, used to treat **hairy cell leukaemia**
- rifampicin, to treat tuberculosis (TB) and other **bacterial infections**
- trimethoprim/sulfamethoxazole, an antibiotic to treat **bacterial infections**
- phenytoin and phenobarbital, to treat **epilepsy**
- oxcarbazepine and carbamazepine, to treat **epilepsy** and **bipolar disorder**
- **St. John's wort** (*Hypericum perforatum*), a herbal remedy to treat **depression**
- **methadone**, used as a **heroin substitute**. Abacavir increases the rate at which methadone is removed from the body. If you are taking methadone, you will be checked for any withdrawal symptoms. Your methadone dose may need to be changed.
- **ribavirin**, for treating **hepatitis C**. Abacavir may make the combination of ribavirin and pegylated interferon less effective at reducing levels of hepatitis C virus in the body.

→ **Tell your doctor or pharmacist** if you are taking any of these. Your doctor may decide to adjust your dose or that you need extra checkups.

## **Pregnancy**

If you are pregnant, think you may be pregnant, or if you are planning to have a baby:

→ **Talk to your doctor** about the risks and benefits of taking Triumeq.

## **Breast-feeding**

**Women who are HIV-positive must not breast-feed** because HIV infection can be passed on to the baby in breast milk.

A small amount of the ingredients in Triumeq can also pass into your breast milk.

If you are breast-feeding, or thinking about breast-feeding:

→ **Talk to your doctor immediately**.

## **Driving and using machines**

**Triumeq can make you dizzy** and have other side effects that make you less alert.

→ **Don't drive or operate machinery** unless you are sure your alertness has not been affected.

### **3. How to take Triumeq**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

- **The usual dose is one tablet once a day**

Swallow the tablet with some liquid. Triumeq can be taken with or without food.

#### **Use in children and adolescents**

Children and adolescents aged between 12 and 17 years and weighing at least 40 kg can take the adult dose of one tablet once a day.

**Do not take an antacid** during the 6 hours before you take Triumeq, or for at least 2 hours after you take it. Other acid-lowering medicines like ranitidine and omeprazole can be taken at the same time as Triumeq.

→ Talk to your doctor for further advice on taking antacid medicines with Triumeq.

**Do not take a calcium or iron supplement** during the 6 hours before you take Triumeq, or for at least 2 hours after you take it.

→ Talk to your doctor for further advice on taking calcium supplements, iron supplements or multivitamins with Triumeq.

#### **If you take more Triumeq than you should**

If you take too many tablets of Triumeq, **contact your doctor or pharmacist for advice**. If possible, show them the Triumeq pack.

#### **If you forget to take Triumeq**

If you miss a dose, take it as soon as you remember. But if your next dose is due within 4 hours, skip the dose you missed and take the next one at the usual time. Then continue your treatment as before.

→ **Don't take a double dose** to make up for a missed dose.

#### **If you have stopped taking Triumeq**

If you have stopped taking Triumeq for any reason — especially because you think you are having side effects, or because you have another illness:

**Talk to your doctor before you start taking it again.** Your doctor will check whether your symptoms were related to a hypersensitivity reaction. If the doctor thinks they may be related to a hypersensitivity reaction, **you will be told never again to take Triumeq, or any other medicine containing abacavir or dolutegravir**. It is important that you follow this advice.

If your doctor advises that you can start taking Triumeq again, you may be asked to take your first doses in a place where you will have ready access to medical care if you need it.

### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, but not everybody gets them.

When you're being treated for HIV, it can be hard to tell whether a symptom is a side effect of Triumeq or other medicines you are taking, or an effect of the HIV disease itself. **So it is very important to talk to your doctor about any changes in your health.**

Even patients who don't have the HLA-B\*5701 gene may still develop **a hypersensitivity reaction** (a serious allergic reaction), described in this leaflet in the panel headed 'Hypersensitivity reactions'. **It is very important that you read and understand the information about this serious reaction.**

**As well as the side effects listed below for Triumeq**, other conditions can develop during combination therapy for HIV.

It is important to read the information in this section under the heading 'Other possible side effects of combination therapy for HIV'.

### **Hypersensitivity Reactions**

Triumeq contains abacavir and dolutegravir. Both of these active ingredients can cause a serious allergic reaction known as a hypersensitivity reaction.

These hypersensitivity reactions have been seen more frequently in people taking medicines that contain abacavir.

#### **Who gets these reactions?**

Anyone taking Triumeq could develop a hypersensitivity reaction, which could be life threatening if they continue to take Triumeq.

You are more likely to develop this reaction if you have a gene called HLA-B\*5701 (but you can get a reaction even if you don't have this gene). You should have been tested for this gene before Triumeq was prescribed for you. If you know you have this gene, tell your doctor.

About 3 to 4 in every 100 patients treated with abacavir in a clinical trial who did not have the HLA-B\*5701 gene developed a hypersensitivity reaction.

#### **What are the symptoms?**

The most common symptoms are:

**fever** (high temperature) and **skin rash**.

Other common symptoms are:

**nausea** (feeling sick), vomiting (being sick), diarrhoea, abdominal (stomach) pain, severe tiredness.

Other symptoms include:

pains in the joints or muscles, swelling of the neck, shortness of breath, sore throat, cough, occasional headaches, inflammation of the eye (conjunctivitis), mouth ulcers, low blood pressure, tingling or numbness of the hands or feet.

### **When do these reactions happen?**

Hypersensitivity reactions can start at any time during treatment with Triumeq, but are more likely during the first 6 weeks of treatment.

### **Contact your doctor immediately:**

- 1 if you get a skin rash, OR**
- 2 if you get symptoms from at least 2 of the following groups:**
  - **fever**
  - **shortness of breath, sore throat or cough**
  - **nausea or vomiting, diarrhoea or abdominal pain**
  - **severe tiredness or aches and pains, or generally feeling ill.**

**Your doctor may advise you to stop taking Triumeq.**

### **If you have stopped taking Triumeq**

If you have stopped taking Triumeq because of a hypersensitivity reaction, **you must NEVER AGAIN take Triumeq, or any other medicine containing abacavir.** If you do, within hours, your blood pressure could fall dangerously low, which could result in death. You should also never again take medicines containing dolutegravir.

If you have stopped taking Triumeq for any reason — especially because you think you are having side effects, or because you have other illness:

**Talk to your doctor before you start again.** Your doctor will check whether your symptoms were related to a hypersensitivity reaction. If the doctor thinks they may have been, **you will then be told never again to take Triumeq, or any other medicine containing abacavir.** You may also be told never again to take any other medicine containing dolutegravir. It is important that you follow this advice.

Occasionally, hypersensitivity reactions have developed in people who start taking abacavir containing products again, but who had only one symptom on the Alert Card before they stopped taking it.

Very rarely, patients who have taken medicines containing abacavir in the past without any symptoms of hypersensitivity have developed a hypersensitivity reaction when they start taking these medicines again.

If your doctor advises that you can start taking Triumeq again, you may be asked to take your first doses in a place where you will have ready access to medical care if you need it.

If you are hypersensitive to Triumeq, return all your unused Triumeq tablets for safe disposal. Ask your doctor or pharmacist for advice.

The Triumeq pack includes an **Alert Card**, to remind you and medical staff about hypersensitivity reactions. **Detach this card and keep it with you at all times.**

## **Very common side effects**

These may affect **more than 1 in 10 people**:

- headache
- diarrhoea
- feeling sick (*nausea*)
- difficulty in sleeping (*insomnia*)
- lack of energy (*fatigue*)

## **Common side effects**

These may affect **up to 1 in 10 people**:

- hypersensitivity reaction (see '*Hypersensitivity reactions*' earlier in this section)
- loss of appetite
- rash
- itching (*pruritus*)
- being sick (*vomiting*)
- stomach (*abdominal*) pain
- stomach (*abdominal*) discomfort
- indigestion
- wind (*flatulence*)
- dizziness
- abnormal dreams
- nightmares
- depression (feelings of deep sadness and unworthiness)
- tiredness
- fever (*high temperature*)
- cough
- irritated or runny nose
- hair loss
- muscle pain and discomfort
- joint pain
- feeling weak
- general feeling of being unwell

Common side effects that may show up in blood tests are:

- an increase in the level of liver enzymes

## **Uncommon side effects**

These may affect **up to 1 in 100 people**:

- inflammation of the liver (*hepatitis*)
- suicidal thoughts and behaviours (particularly in patients who have had depression or mental health problems before)

Uncommon side effects that may show up in blood tests are:

- a decreased number of cells involved in blood clotting (*thrombocytopenia*).
- a low red blood cell count (*anaemia*) or low white blood cell count (*neutropenia*)
- an increase in sugar (*glucose*) in the blood
- an increase in triglycerides (type of fat) in the blood

## **Rare side effects**

These may affect **up to 1 in 1000 people**:

- inflammation of the pancreas (*pancreatitis*)

- breakdown of muscle tissue.

Rare side effects that may show up in blood tests are:

- increase in an enzyme called *amylase*

### **Very rare side effects**

These may affect **up to 1 in 10,000** people:

- numbness, tingly feelings in the skin (pins and needles)
- sensation of weakness in the limbs
- skin rash, which may form blisters and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge) (*erythema multiforme*)
- a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (*Stevens–Johnson syndrome*), and a more severe form causing skin peeling in more than 30% of the body surface (*toxic epidermal necrolysis*).

Very rare side effects that may show up in blood tests are:

- a failure of the bone marrow to produce new red blood cells (*pure red cell aplasia*).

If you get any side effects

→ **Talk to your doctor.** This includes any possible side effects not listed in this leaflet.

### **Other possible side effects of combination therapy for HIV**

Combination therapy such as Triumeq may cause other conditions to develop during HIV treatment.

#### **Symptoms of infection and inflammation**

People with advanced HIV infection or AIDS have weak immune systems, and are more likely to develop serious infections (*opportunistic infections*). Such infections may have been “silent” and not detected by the weak immune system before treatment was started. After starting treatment, the immune system becomes stronger, and may attack the infections, which can cause symptoms of infection or inflammation. Symptoms usually include **fever**, plus some of the following:

- headache
- stomach ache
- difficulty breathing

In rare cases, as the immune system becomes stronger, it can also attack healthy body tissue (*autoimmune disorders*). The symptoms of autoimmune disorders may develop many months after you start taking medicine to treat your HIV infection. Symptoms may include:

- palpitations (rapid or irregular heartbeat) or tremor
- hyperactivity (excessive restlessness and movement)
- weakness beginning in the hands and feet and moving up towards the trunk of the body.

**If you get any symptoms of infection** and inflammation or if you notice any of the symptoms above:

→ **Tell your doctor immediately.** Don’t take other medicines for the infection without your doctor’s advice.

#### **Your body shape may change**

People taking combination therapy for HIV may find that their body shape changes, because of changes in fat distribution:

- Fat may be lost from the legs, arms or face.
- Extra fat may build up around the tummy (abdomen), or on the breasts or internal organs.
- Fatty lumps (sometimes called buffalo hump) may appear on the back of the neck.

If you notice changes in your body shape:

→ **Tell your doctor.**

## **Lactic acidosis is a rare but serious side effect**

Some people taking Triumeq, or other medicines like it (NRTIs), develop a condition called lactic acidosis, together with an enlarged liver.

Lactic acidosis is caused by a build-up of lactic acid in the body. It is rare; if it happens, it usually develops after a few months of treatment. It can be life-threatening, causing failure of internal organs.

Lactic acidosis is more likely to develop in people who have liver disease, or in obese (very overweight) people, especially women.

### **Signs of lactic acidosis include:**

- deep, rapid, difficult breathing
- drowsiness
- numbness or weakness in the limbs
- feeling sick (*nausea*), being sick (*vomiting*)
- stomach pain.

During your treatment, your doctor will monitor you for signs of lactic acidosis. If you have any of the symptoms listed above or any other symptoms that worry you:

**See your doctor as soon as possible.**

## **Joint pain, stiffness and bone problems**

Some people taking combination therapy for HIV develop a condition called *osteonecrosis*. In this condition, parts of the bone tissue die because of reduced blood supply to the bone. People may be more likely to get this condition:

- if they have been taking combination therapy for a long time
- if they are also taking anti-inflammatory medicines called corticosteroids
- if they drink alcohol
- if their immune systems are very weak
- if they are overweight.

### **Signs of osteonecrosis include:**

- stiffness in the joints
- aches and pains (especially in the hip, knee or shoulder)
- difficulty moving.

If you notice any of these symptoms:

**→ Tell your doctor.**

## **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Triumeq**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and bottle after EXP.

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

This medicine does not require any special temperature storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Triumeq contains

- The active substances are dolutegravir, abacavir and lamivudine. Each tablet contains dolutegravir sodium equivalent to 50 mg dolutegravir, 600 mg abacavir (as sulfate) and 300 mg lamivudine.
- The other ingredients are mannitol (E421), microcrystalline cellulose, povidone K29/32, sodium starch glycollate, magnesium stearate. Opadry II Purple 85F90057 (containing polyvinyl alcohol – part hydrolyzed, titanium dioxide, macrogol/PEG, talc, iron oxide black and iron oxide red).

### What Triumeq looks like and contents of the pack

Triumeq film-coated tablets are purple, biconvex, oval tablets, debossed with "572 Tri" on one side. The film-coated tablets are provided in bottles containing 30 tablets.

The bottle contains a desiccant to reduce moisture. Once the bottle has been opened keep the desiccant in the bottle, do not remove it.

Multipacks containing 90 film-coated tablets (3 packs of 30 film-coated tablets) are also available. Not all pack sizes may be available in your country.

### Marketing Authorisation Holder

ViiV Healthcare UK Limited, 980 Great West Road, Brentford, Middlesex, TW8 9GS, United Kingdom.

### Manufacturer

Glaxo Wellcome, S.A., Avda. Extremadura 3, 09400 Aranda De Duero, Burgos, Spain

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### België/Belgique/Belgien

ViiV Healthcare sprl/bvba  
Tél/Tel: + 32 (0) 10 85 65 00

#### Lietuva

GlaxoSmithKline Lietuva UAB  
Tel: + 370 5 264 90 00  
info.lt@gsk.com

#### България

ГлаксоСмитКлейн ЕООД  
Тел.: + 359 2 953 10 34

#### Luxembourg/Luxemburg

ViiV Healthcare sprl/bvba  
Belgique/Belgien  
Tél/Tel: + 32 (0) 10 85 65 00

#### Česká republika

GlaxoSmithKline, s.r.o.  
Tel: + 420 222 001 111  
cz.info@gsk.com

#### Magyarország

GlaxoSmithKline Kft.  
Tel.: + 36 1 225 5300

#### Danmark

GlaxoSmithKline Pharma A/S  
Tlf: + 45 36 35 91 00  
[dk-info@gsk.com](mailto:dk-info@gsk.com)

#### Malta

GlaxoSmithKline (Malta) Limited  
Tel: + 356 21 238131

**Deutschland**

ViiV Healthcare GmbH  
Tel.: + 49 (0)89 203 0038-10  
[vii.med.info@viivhealthcare.com](mailto:vii.med.info@viivhealthcare.com)

**Eesti**

GlaxoSmithKline Eesti OÜ  
Tel: + 372 6676 900  
[estonia@gsk.com](mailto:estonia@gsk.com)

**Ελλάδα**

GlaxoSmithKline A.E.B.E.  
Τηλ: + 30 210 68 82 100

**España**

Laboratorios ViiV Healthcare, S.L.  
Tel: + 34 902 051 260  
[es-ci@viivhealthcare.com](mailto:es-ci@viivhealthcare.com)

**France**

ViiV Healthcare SAS  
Tél.: + 33 (0)1 39 17 69 69  
[Infomed@viivhealthcare.com](mailto:Infomed@viivhealthcare.com)

**Hrvatska**

GlaxoSmithKline d.o.o.  
Tel: + 385 1 6051 999

**Ireland**

GlaxoSmithKline (Ireland) Limited  
Tel: + 353 (0)1 4955000

**Ísland**

GlaxoSmithKline ehf.  
Sími: + 354 530 3700

**Italia**

ViiV Healthcare S.r.l  
Tel: + 39 (0)45 9212611

**Κύπρος**

GlaxoSmithKline (Cyprus) Ltd  
Τηλ: + 357 22 39 70 00  
[gskcyprus@gsk.com](mailto:gskcyprus@gsk.com)

**Latvija**

GlaxoSmithKline Latvia SIA  
Tel: + 371 67312687

**Nederland**

ViiV Healthcare BV  
Tel: + 31 (0)30 6986060  
[contact-nl@viivhealthcare.com](mailto:contact-nl@viivhealthcare.com)

**Norge**

GlaxoSmithKline AS  
Tlf: + 47 22 70 20 00  
[firmapost@gsk.no](mailto:firmapost@gsk.no)

**Österreich**

GlaxoSmithKline Pharma GmbH  
Tel: + 43 (0)1 97075 0  
[at.info@gsk.com](mailto:at.info@gsk.com)

**Polska**

GSK Services Sp. z o.o.  
Tel.: + 48 (0)22 576 9000

**Portugal**

VIIHV HIV HEALTHCARE, UNIPESSOAL, LDA  
Tel: + 351 21 094 08 01  
[vii.vi.pt@viivhealthcare.com](mailto:vii.vi.pt@viivhealthcare.com)

**România**

GlaxoSmithKline (GSK) S.R.L.  
Tel: + 4021 3028 208

**Slovenija**

GlaxoSmithKline d.o.o.  
Tel: + 386 (0)1 280 25 00  
[medical.x.si@gsk.com](mailto:medical.x.si@gsk.com)

**Slovenská republika**

GlaxoSmithKline Slovakia s. r. o.  
Tel: + 421 (0)2 48 26 11 11  
[recepacia.sk@gsk.com](mailto:recepacia.sk@gsk.com)

**Suomi/Finland**

GlaxoSmithKline Oy  
Puh/Tel: + 358 (0)10 30 30 30  
[Finland.tuoteinfo@gsk.com](mailto:Finland.tuoteinfo@gsk.com)

**Sverige**

GlaxoSmithKline AB  
Tel: + 46 (0)8 638 93 00  
[info.produkt@gsk.com](mailto:info.produkt@gsk.com)

**United Kingdom**

ViiV Healthcare UK Limited  
Tel: + 44 (0)800 221441

lv-epasts@gsk.com

customercontactuk@gsk.com

**This leaflet was last revised in {month YYYY}.**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.